

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
11 December 2003 (11.12.2003)

PCT

(10) International Publication Number  
**WO 03/101375 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K**
- (21) International Application Number: PCT/EP03/05691
- (22) International Filing Date: 30 May 2003 (30.05.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
2,388,049 30 May 2002 (30.05.2002) CA
- (71) Applicant (for all designated States except US): **IMMUNOTECH S.A.** [AR/AR]; Florida 1, 10th Floor, Office 36, Ciudad Autonoma de Buenos Aires (AR).

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **LOPEZ, Ricardo, Augustin** [IT/AR]; Camarones 2664, Ciudad Autonoma de Buenos Aires (AR).

- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— of inventorship (Rule 4.17(iv)) for US only

**Published:**



— without international search report and to be republished upon receipt of that report

- (74) Agents: **PISTOLESI, Roberto** et al.; Dragotti & Associati Srl, Galleria San Babila 4/C, I-20122 Milano (IT).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

- (54) Title: **IMMUNOSTIMULATORY OLIGONUCLEOTIDES AND USES THEREOF**



- (57) Abstract: Oligonucleotides containing the non-palindromic sequence motif  $X_1X_2X_3X_4X_5X_6X_7X_8$ , wherein  $X_1$  is C or T,  $X_2$  is C, T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are Ts and wherein C never precedes G (in other terms, the nucleic acid motif does not consist of a CpG oligonucleotides) that have the ability to stimulate the immune response of animals of the order *Primate*, including humans are disclosed. The stimulation of the immune response is characterized by stimulation of proliferation, differentiation, cytokine production and antibody production in B-cells and cell differentiation and cytokine production in plasmacytoid dendritic cells.

## IMMUNOSTIMULATORY OLIGONUCLEOTIDES AND USES THEREOF

### FIELD OF THE INVENTION

The present invention refers especially to oligonucleotides containing the non-palindromic sequence motif:

5  $X_1X_2X_3X_4X_5X_6X_7X_8$ ,

wherein  $X_1$  is C or T,  $X_2$  is C, T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are Ts and wherein C never precedes G. These oligonucleotides were shown to be immunostimulatory in animals of the order *Primate*, including  
10 humans.

### RELEVANT REFERENCES

---

#### PATENT DOCUMENTS

15 US 5,663,153; US 5,723,335; US 6,090,791; US 6,194,388; US 6,207,646;  
US 6,239,116; US 6,429,199; US 6,544,518; US 6,514,948; US 6,498,148;  
US 6,429,199; US 6,426,336; US 6,406,705; US 6,455,689; US 20010044416;  
US 20020156033; US 20020165178; US 20020198165; US 2002137714;  
US 20030050268; US 2003040499; EP 0468520; WO 95/26204; WO  
20 96/02555; WO 98/40100; WO 98/52962; WO 99/33488; WO 99/56755; WO  
00/62802; WO 01/00231; WO 01/00232; WO 01/22972; WO 01/55370; WO  
00/014217; WO 00/061151; WO 00/067023; WO 00/067787; WO 00/075304;  
WO 01/002007; WO 01/007055; WO 01/022990; WO 01/048018; WO  
01/055341; WO 01/062909; WO 01/062910; WO 01/068077; WO 01/068116;  
25 WO 01/068143; WO 01/068144; WO 01/083503; WO 01/093905; WO

02/026757; WO 02/056909; WO 02/069369; WO 02/095027; WO 02/102825;  
WO 03/002065; WO 94/25588; WO 99/56755; WO 99/62923.

#### OTHER REFERENCES

- 5 Ahmad-Nejad P et al., Bacterial CpG-DNA and lipopolysaccharides activate Toll-like receptors at distinct cellular compartments. *Eur. J. Immunol.* 32:1958-1968, (2002).
- Angier N. Microbe DNA Seen as Alien By Immune System, *New York Times*, 11, (1995).
- 10 Auf G et al., CpG-oligodeoxynucleotide rejection of a neuroblastoma in A/J mice does not induce a paraneoplastic disease. *Neurosci. Lett.* 327:189-192, (2002).
- Ayash-Rashkovsky M et al., Generation of Th1 immune responses to inactivated, gp120-depleted HIV-1 in mice with a dominant Th2 biased  
15 immune profile via immunostimulatory oligonucleotides--relevance to AIDS vaccines in developing countries. *Vaccine.* 20:2684-2692, (2002).
- Azad RF et al., Antiviral Activity of a Phosphorothioate Oligonucleotide Complementary to RNA of the Human Cytomegalovirus Major Immediate-Early Region, *Antimicrobial Agents and Chemotherapy*, 37:1945-1954,  
20 (1993).
- Azuma, Biochemical and Immunological Studies on Cellular Components of Tubercle Bacilli. *Kekkaku*, 69. 9:45-55, (1992).

Ballas ZK et al., Induction of NK activity in murine and human cells by CpG motifs in oligodeoxynucleotides and bacterial DNA. *J. Immunol.* 157, 5:1840-1845, (1996).

Bauer M et al., Bacterial CpG-DNA triggers activation and maturation of human CD11c-, CD123+ dendritic cells. *J. Immunol.* 166:5000-5007 (2001).

---

Bauer S et al., H. Bacterial CpG-DNA licenses TLR9. *Curr. Top. Microbiol. Immunol.* 270:145-154 (2002).

Bauer S et al., Human TLR9 confers responsiveness to bacterial DNA via species-specific CpG motif recognition. *Proc. Natl. Acad. Sci. USA*, 98: 9237-9242, (2001).

Beignon AS et al., Immunization onto bare skin with synthetic peptides: immunomodulation with a CpG-containing oligodeoxynucleotide and effective priming of influenza virus-specific CD4+ T cells. *Immunology*, 105:204-2012 (2002).

Boggs RT et al., Characterization and modulation of immune stimulation by modified oligonucleotides. *Antisense Nucleic Acid Drug Dev.* 7, 5:461-471, (1997).

Bohle B. CpG motifs as possible adjuvants for the treatment of allergic diseases. *Int. Arch. Allergy Immunol.* 129:198-203, (2002).

Bozza S et al., Vaccination of mice against invasive aspergillosis with recombinant *Aspergillus* proteins and CpG oligodeoxynucleotides as adjuvants. *Microbes Infect.* 4:1281-1290, (2002).

Branda RF et al., Amplification of antibody production by phosphorothioate oligodeoxynucleotides. *J. Lab. Clin. Med.* 128, 3:329-338, (1996).

Carlson GA. Postexposure prophylaxis against transmissible spongiform encephalopathies: CpG oligodeoxynucleotides in mice. *Lancet*, 360:184,  
5 (2002).

Carpentier et al., Oligodeoxynucleotides containing CpG motifs can induce rejection of a neuroblastoma in mice. *Cancer Res.* 59, 5429-5432, (1999).

Chace JH et al., Bacterial DNA-induced NK cell IFN $\gamma$  production is dependent on macrophage secretion of IL-12. *Clin. Immunol. Immunopathol.* 84, 185-  
10 193, (1997).

Chen W et al., Enhancement of antigen-presenting ability of B lymphoma cells by immunostimulatory CpG-oligonucleotides and anti-CD40 antibody. *Immunol. Lett.* 77:17-23 (2001).

Choudhury BK et al., In vivo role of p38 mitogen-activated protein kinase in  
15 mediating the anti-inflammatory effects of CpG oligodeoxynucleotide in murine asthma. *J. Immunol.* 169:5955-5961, (2002).

Chu RS et al., CpG oligodeoxynucleotides act as adjuvants that switch on T helper 1 (Th1) immunity. *J. Exp. Med.* 186, 1623-1631, (1997).

Chuang TH et al., Toll-like receptor 9 mediates CpG-DNA signaling. *J.*  
20 *Leukoc. Biol.* 71:538-544, (2002).

Cowdery JS et al., Bacterial DNA induces NK cells to produce IFN-gamma in vivo and increases the toxicity of lipopolysaccharides. *J. Immunol.* 156, 12:4570-4575, (1996).

5 Dalpke AH et al., Phosphodiester CpG oligonucleotides as adjuvants: polyguanosine runs enhance cellular uptake and improve immunostimulative activity of phosphodiester CpG oligonucleotides in vitro and in vivo. *Immunology*, 106:102-1012, (2002).

Davis HL et al., CpG DNA is a potent enhancer of specific immunity in mice immunized with recombinant hepatitis B surface antigen. *J. Immunol.* 160, 10 870-876, (1998).

Decker T et al., Effect of immunostimulatory CpG-oligonucleotides in chronic lymphocytic leukemia B cells. *Leuk. Lymphoma.* 42:301-307 (2001).

Decker T et al., Immunostimulatory CpG-oligonucleotides cause proliferation, cytokine production, and an immunogenic phenotype in chronic lymphocytic leukemia B cells. *Blood*, 95:999-1006 (2000). 15

Decker T et al., Peschel C. Immunostimulatory CpG-oligonucleotides induce functional high affinity IL-2 receptors on B-CLL cells: costimulation with IL-2 results in a highly immunogenic phenotype. *Exp. Hematol.* 28:558-68 (2000).

Decker T et al., Sensitization of B-cell chronic lymphocytic leukemia cells to recombinant immunotoxin by immunostimulatory phosphorothioate oligodeoxynucleotides. *Blood*, 99:1320-1326, (2002). 20

Diwan M et al., Enhancement of immune responses by co-delivery of a CpG oligodeoxynucleotide and tetanus toxoid in biodegradable nanospheres. *J. Control Release*. 85:247-62, (2002).

Dumais N et al., Mucosal immunization with inactivated human immunodeficiency virus plus CpG oligodeoxynucleotides induces genital immune responses and protection against intravaginal challenge. *J. Infect. Dis.* 186:1098-105 (2002).

Erb KJ et al., Infection of mice with *Mycobacterium bovis*-*Bacillus Calmette-Guerin* (BCG) suppresses allergen-induced airway eosinophilia. *J. Exp. Med.* 187, 4:561-569, (1998).

Fattorini L et al., Recombinant GroES in combination with CpG oligodeoxynucleotides protects mice against *Mycobacterium avium* infection. *J. Med. Microbiol.* 51:1071-1079, (2002).

Gierynska M et al., Induction of CD8 T-cell-specific systemic and mucosal immunity against herpes simplex virus with CpG-peptide complexes. *J. Virol.* 76:6568-6576 (2002).

Gill RF et al., Enhancement of rat tear IgA antibody responses following intranasal immunization with antigen and CpG ODN. *Curr. Eye Res.* 24:228-233, (2002).

Gursel M et al., CpG oligodeoxynucleotides induce human monocytes to mature into functional dendritic cells. *Eur. J. Immunol.* 32:2617-2622 (2002).

Gursel M et al., Differential and competitive activation of human immune cells by distinct classes of CpG oligodeoxynucleotide. *J. Leukoc. Biol.* 71:813-820, (2002).

Hadden J et al., Immunopharmacology, JAMA, 268, 20:2964-2969, (1992).

5 Hadden J et al., Immunostimulants. *TIPS*, 141:169-174, (1993).

---

Halpern MD et al., Bacterial DNA induces murine interferon-gamma production by stimulation of interleukin-12 and tumor necrosis factor-alpha. *Cell Immunol.* 167, 1:72-78, (1996).

Harandi AM et al., A protective role of locally administered immunostimulatory  
10 CpG oligodeoxynucleotide in a mouse model of genital herpes infection. *J. Virol.* 77:953-962 (2003).

Hartmann et al., Mechanism and function of a newly identified CpG DNA motif in human primary B cells. *J. Immunol.* 164, 944-952, (2000).

Hartmann G et al., Delineation of a CpG phosphorothioate  
15 oligodeoxynucleotide for activating primate immune responses in vitro and in vivo. *J. Immunol.* 164, 1617-1624, (2000).

Heckelsmiller K et al., Combined dendritic cell- and CpG oligonucleotide-based immune therapy cures large murine tumors that resist chemotherapy. *Eur. J. Immunol.* 32:3235-45, (2002).

20 Heckelsmiller K et al., Peritumoral CpG DNA elicits a coordinated response of CD8 T cells and innate effectors to cure established tumors in a murine colon carcinoma model. *J. Immunol.* 169:3892-3899, (2002).



- Horner AA et al., Immunostimulatory sequence oligodeoxynucleotide-based vaccination and immunomodulation: two unique but complementary strategies for the treatment of allergic diseases. *J. Allergy Clin. Immunol.* 110:706-712 (2002).
- 5 Horner AA et al., Optimized conjugation ratios lead to allergen immunostimulatory oligodeoxynucleotide conjugates with retained immunogenicity and minimal anaphylactogenicity. *J. Allergy Clin. Immunol.* 110:413-420, (2002).
- 10 Hornung V et al., Quantitative expression of toll-like receptor 1-10 mRNA in cellular subsets of human peripheral blood mononuclear cells and sensitivity to CpG oligodeoxynucleotides. *J. Immunol.* 168:4531-4537, (2002).
- Hussain I et al., Modulation of murine allergic rhinosinusitis by CpG oligodeoxynucleotides. *Laryngoscope*, 112:1819-1826, (2002).
- 15 Ioannou XP et al., CpG-containing oligodeoxynucleotides, in combination with conventional adjuvants, enhance the magnitude and change the bias of the immune responses to a herpesvirus glycoprotein. *Vaccine* 21:127-37, (2002).
- Ioannou XP et al., The immunogenicity and protective efficacy of bovine herpesvirus 1 glycoprotein D plus Emulsigen are increased by formulation with CpG oligodeoxynucleotides. *J. Virol.* 76:9002-9010, (2002).
- 20 Ishii KJ, et al., Potential role of phosphatidylinositol 3 kinase, rather than DNA-dependent protein kinase, in CpG DNA-induced immune activation. *J. Exp. Med.* 196:269-274 (2002).

Jahrsdorfer B et al., CpG DNA increases primary malignant B cell expression of costimulatory molecules and target antigens. *J. Leukoc. Biol.* 69:81-88 (2001).

Jahrsdorfer B et al., Modulation of malignant B cell activation and apoptosis  
5 by bcl-2 antisense ODN and immunostimulatory CpG ODN. *J. Leukoc. Biol.*  
72:83-92, (2002).

Jain VV et al., CpG-oligodeoxynucleotides inhibit airway remodeling in a murine model of chronic asthma. *J. Allergy Clin. Immunol.* 110:867-872, (2002).

10 Jiang W et al., Mucosal immunization with helicobacter, CpG DNA, and cholera toxin is protective. *Infect. Immun.* 71:40-46, (2003).

Joseph A et al., Liposomal immunostimulatory DNA sequence (ISS-ODN): an efficient parenteral and mucosal adjuvant for influenza and hepatitis B vaccines. *Vaccine.* 20:3342-3354 (2002).

15 Juffermans NP et al., CpG oligodeoxynucleotides enhance host defense during murine tuberculosis. *Infect. Immun.* 70:147-152 (2002).

Jung J et al., Distinct response of human B cell subpopulations in recognition of an innate immune signal, CpG DNA. *J. Immunol.* 169:2368-2373, (2002).

Kadowaki N et al., Distinct CpG DNA and polyinosinic-polycytidylic acid  
20 double-stranded RNA, respectively, stimulate CD11c<sup>-</sup> type 2 dendritic cell precursors and CD11c<sup>+</sup> dendritic cells to produce type I IFN. *J. Immunol.*, 166, 2291- 2295, (2001).

Kadowaki N et al., Subsets of human dendritic cell precursors express different toll-like receptors and respond to different microbial antigens. *J. Exp. Med.* 17;194:863-869 (2001).

Kataoka T et al., Antitumor Activity of Synthetic Oligonucleotides with Sequences from cDNA Encoding Proteins of Mycobacterium bovis BCG. *Jpn. J. Cancer Res.*, 83:244-247, (1992).

Kimura T et al., Binding of Oligoguanylate to Scavenger Receptors Is Required for Oligonucleotides to Augment NK Cell Activity and Induce IFN. *J. Biochem.* 116, 5:991-994, (1994).

Kitagaki K et al., Immunomodulatory effects of CpG oligodeoxynucleotides on established th2 responses. *Clin. Diagn. Lab. Immunol.* 9:1260-1269, (2002).

Kline JN et al., CpG oligonucleotides can reverse as well as prevent TH2-mediated inflammation in a murine model of asthma. *J. Invest. Med.*, 45, 7:298A, (1997).

Kline JN et al., Immune redirection by CpG oligonucleotides. Conversion of a Th2 response to a Th1 response in a murine model of asthma. *J. Invest. Med.*, 45, 3:282A, (1997).

Kline JN et al., Modulation of airway inflammation by CpG oligodeoxynucleotides in a murine model of asthma. *J. immunol.* 15, 2555-2559, (1998).

Kline JN et al., Treatment of established asthma in a murine model using CpG oligodeoxynucleotides. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 283:170-179, (2002).

Klinman DM et al., CpG motifs present in bacteria DNA rapidly induce  
5 lymphocytes to secrete interleukin 6, interleukin 12, and interferon gamma.

---

*Proc. Natl. Acad. Sci. USA*, 93, 7:2879-2883, (1996).

Krieg AM et al., A Role for Endogenous Retroviral Sequences in the Regulation of Lymphocyte Activation. *J. Immunol.*, 143, 2448-2451, (1989).

Krieg AM et al, Phosphorothioate Oligodeoxynucleotides: Antisense or Anti-Protein?, *Antisense Res. Develop.* 5:241, (1995).  
10

Krieg AM et al. Leukocyte Stimulation by Oligodeoxynucleotides. *Applied Antisense Oligonucleotide Technology*, 431-448, (1998).

Krieg AM et al., CpG motifs in bacterial DNA trigger direct B-cell activation. *Nation*, 374:546-549, (1995).

15 Krieg AM et al., Oligodeoxynucleotide modifications determine the magnitude of B cell stimulation by CpG motifs. *Antisense Nucleic Acid Drug Dev.*, 6, 2:133-139, (1996).

Krieg AM et al., The role of CpG dinucleotides in DNA vaccines. *Trends in Microbiology*, 6, 23-37, (1998).

20 Krieg AM, An innate immune defense mechanism based on the recognition of CpG motifs in microbial DNA. *J. Lab. Clin. Med.*, 128, 2:128-33, (1996).

Krieg AM. CpG motifs in bacterial DNA and their immune effects. *Annu. Rev. Immunol.* 20:709-760 (2002).

Krieg AM. GpG motifs in bacterial DNA and their immune effects. *Annu. Rev. Immunol.*, 20, 709-760, (2002).

- 5 Krug A et al., Identification of CpG oligonucleotide sequences with high induction of IFN-alpha/beta in plasmacytoid dendritic cells. *Eur. J. Immunol.* 31:2154-2163 (2001).

- Krug A et al., Toll-like receptor expression reveals CpG DNA as a unique microbial stimulus for plasmacytoid dendritic cells which synergizes with  
10 CD40 ligand to induce high amounts of IL-12. *Eur. J. Immunol.* 31:3026-3037, (2002).

Kuramoto, et al., Oligonucleotide Sequences Required for Natural Killer Cell Activation. *Jpn. J. Cancer Res.* 83:1128-1131, (1992).

- Liang H et al., Activation of human B cells by phosphorothioate  
15 oligodeoxynucleotides. *J. Clin. Invest.* 98, 1119-1129, (1996).

Lingnau K et al., Poly-L-arginine synergizes with oligodeoxynucleotides containing CpG-motifs (CpG-ODN) for enhanced and prolonged immune responses and prevents the CpG-ODN-induced systemic release of pro-inflammatory cytokines. *Vaccine.* 20:3498-3508, (2002).

- 20 Lipford GB et al., CpG-containing synthetic oligonucleotides promote B and cytotoxic T cell responses to protein antigen: a new class of vaccine adjuvants. *Eur. J. Immunol.*, 27:2340-2344, (1997).

Lipford GB et al., Immunostimulatory DNA: sequence-dependent production of potentially harmful or useful cytokines. *Eur. J. Immunol.* 27:3420-3426, (1997).

Macfarlane DE and Manzel L, Antagonism of immunostimulatory CpG-oligodeoxynucleotides by quinacrine, chloroquine, and structurally related compounds. *J. Immunol.*, 160, 3:1122-1131, (1998).

Magone MT et al., Systemic or mucosal administration of immunostimulatory DNA inhibits early and late phases of murine allergic conjunctivitis. *Eur. J. Immunol.* 30, 1841-1850, (2000).

10 Mariotti S et al., Immunogenicity of anti-Haemophilus influenzae type b CRM197 conjugate following mucosal vaccination with oligodeoxynucleotide containing immunostimulatory sequences as adjuvant. *Vaccine.* 20:2229-2239, (2002).

Matson S et al., Nonspecific suppression of [3H]thymidine incorporation by "control" oligonucleotides. *Antisense Res. Dev.*, 2, 4:325-330, (1992).

Maurer T et al., CpG-DNA aided cross-presentation of soluble antigens by dendritic cells. *Eur. J. Immunol.* 32:2356-2364.

McCluskie MJ et al., Parenteral and mucosal prime-boost immunization strategies in mice with hepatitis B surface antigen and CpG DNA. *FEMS Immunol. Med. Microbiol.* 32:179-185, (2002).

Messina et al., Stimulation of in vitro Murine Lymphocyte Proliferation by Bacterial DNA. *J. Immunol.*, 147, 6:1759-1764, (1991).

Messina, et al., The Influence of DNA Structure on the in vitro Stimulation of Murine Lymphocytes by Natural and Synthetic Polynucleotide Antigens. *Cell. Immunol.*, 147:148-157, (1993).

Miconnet I et al., CpG are efficient adjuvants for specific CTL induction against tumor antigen-derived peptide. *J. Immunol.* 168:1212-1218 (2002).

Mojcik C et al., Administration of a Phosphorothioate Oligonucleotide Antisense Murine Endogenous Retroviral MCF env Causes Immune Effect in vivo in a Sequence-Specific Manner. *Clin. Immunol. Immunopathology.* 67, 2:130-136, (1993).

10 Moldovenu GJ et al., CpG DNA, a novel immune enhancer for systemic and mucosal immunization with influenza virus. *Vaccine.* 16, 1216-1224, (1998).

O'Hagan DT et al. Synergistic adjuvant activity of immunostimulatory DNA and oil/water emulsions for immunization with HIV p55 gag antigen. *Vaccine.* 20:3389-98, (2002).

15 Olbrich AR et al., Effective postexposure treatment of retrovirus-induced disease with immunostimulatory DNA containing CpG motifs. *J. Virol.* 76:11397-404, (2002).

Pal S et al., Immunization with the *Chlamydia trachomatis* mouse pneumonitis major outer membrane protein by use of CpG oligodeoxynucleotides as an adjuvant induces a protective immune response against an intranasal chlamydial challenge. *Infect. Immun.* 70:4812-4817, (2002).

20

Park Y et al., Cutting Edge: CpG DNA inhibits dendritic cell apoptosis by up-regulating cellular inhibitor of apoptosis proteins through the phosphatidylinositol-3'-OH kinase pathway. *J. Immunol.* 168:5-8 (2002).

Pisetsky et al., Stimulation of Murine Lymphocyte Proliferation by a  
5 Phosphorothioate Oligonucleotide with Antisense Activity for Herpes Simplex  
Virus. *Life Science*, 54, 101-107 (1994).

Pisetsky, Immunological Consequences of Nucleic Acid Therapy. *Antisense Res. Development*, 5:219-225 (1995).

Pisetsky, The Immunological Properties of DNA. *J. Immunol.*, 421-423  
10 (1996).

Rachmilewitz D et al., Immunostimulatory DNA ameliorates experimental and spontaneous murine colitis. *Gastroenterology*. 122:1428-1441, (2002).

Rankin R et al., CpG motif identification for veterinary and laboratory species demonstrates that sequence recognition is highly conserved. *Antisense*  
15 *Nucleic Acid Drug Dev.*, 11:333-340 (2001).

Rankin R et al., CpG-containing oligodeoxynucleotides augment and switch the immune responses of cattle to bovine herpesvirus-1 glycoprotein D. *Vaccine*, 20:3014-3022, (2002).

Rhee EG et al., Vaccination with heat-killed leishmania antigen or  
20 recombinant leishmanial protein and CpG oligodeoxynucleotides induces long-term memory CD4+ and CD8+ T cell responses and protection against leishmania major infection. *J. Exp. Med.* 195:1565-1573 (2002).



Rothenfusser S et al., Plasmacytoid dendritic cells: the key to CpG. *Human Immunol.* 63:1111-1119, (2002).

Santeliz JV et al., Amb a 1-linked CpG oligodeoxynucleotides reverse established airway hyperresponsiveness in a murine model of asthma. *J. Allergy Clin. Immunol.* 109:455-462, (2002).

Sato et al., Immunostimulatory DNA Sequences Necessary for Effective Intradermal Gene Immunization. *Science*, 273:352-354, (1996).

Schultz KR et al., Chloroquine prevention of murine MHC-disparate acute graft-versus-host disease correlates with inhibition of splenic response to CpG oligodeoxynucleotides and alterations in T-cell cytokine production. *Biol. Blood Marrow Transplant*, 8:648-655 (2002).

Schwartz DA et al., CpG motifs in bacterial DNA cause inflammation in the lower respiratory tract. *J. Clin. Invest.*, 100, 1:68-73, (1997).

Senuma A et al., Therapeutic effect of CpG motifs on the development of chronic graft-versus-host disease in mice. *Cytokine*. 20:23-29, (2002).

Sethi S et al., Postexposure prophylaxis against prion disease with a stimulator of innate immunity. *Lancet*, 360:229-230, (2002).

Sfondrini L et al., Absence of the CD1 molecule up-regulates antitumor activity induced by CpG oligodeoxynucleotides in mice. *J. Immunol.* 169:151-158, (2002).

- Shirota H et al., B cells capturing antigen conjugated with CpG oligodeoxynucleotides induce Th1 cells by elaborating IL-12. *J. Immunol.* 169:787-94, (2002).
- Sparwasser T et al., Bacterial DNA and immunostimulatory CpG  
5 oligodeoxynucleotides trigger maturation and activation of murine dendritic cells. *Eur. J. Immunol.* 28, 2045-2054, 1998.
- Stern BV et al., Vaccination with tumor peptide in CpG adjuvant protects via IFN-gamma-dependent CD4 cell immunity. *J. Immunol.* 168:6099-6105, (2002).
- 10 Sweet MJ et al., Colony-stimulating factor-1 suppresses responses to CpG DNA and expression of toll-like receptor 9 but enhances responses to lipopolysaccharide in murine macrophages. *J. Immunol.* 168:392-399 (2002).
- Takauji R et al., CpG-DNA-induced IFN-alpha production involves p38 MAPK-dependent STAT1 phosphorylation in human plasmacytoid dendritic  
15 cell precursors. *J. Leukoc. Biol.* 72:1011-1019 (2002).
- Takeshita F et al., Cutting edge: Role of Toll-like receptor 9 in CpG DNA-induced activation of human cells. *J. Immunol.* 67:3555-3558, (2001).
- Tokunaga T et al. Synthetic Oligonucleotides with Particular Base Sequences from the cDNA Encoding Proteins of *Mycobacterium bovis* BCG Induce  
20 Interferons and Activate Natural Killer Cells. *Microbiol. Immunol.*, 36, 1:55-66, (1992).

Tokunaga T et al., A synthetic single-stranded DNA, poly(dG,dC), induces interferon-alpha/beta and -gamma, augments natural killer activity, and suppresses tumor growth. *Jpn. J. Cancer Res.*, 79:682-686, (1988).

Van der Stede Y et al., CpG-oligodinucleotides as an effective adjuvant in  
5 pigs for intramuscular immunizations. *Vet. Immunol. Immunopathol.* 86:31-41,  
(2002).

Verthelyi D et al., CpG oligodeoxynucleotides as vaccine adjuvants in  
primates. *J. Immunol.* 168:1659-1663 (2002).

Verthelyi D et al., Human peripheral blood cells differentially recognize and  
10 respond to two distinct CPG motifs. *J. Immunol.* 166: 2372-2377, (2001).

Vicari AP et al., Reversal of tumor-induced dendritic cell paralysis by CpG  
immunostimulatory oligonucleotide and anti-interleukin 10 receptor antibody.  
*J. Exp. Med.* 196:541-549, (2002).

Vollmer J et al., Highly immunostimulatory CpG-free oligodeoxynucleotides  
15 for activation of human leukocytes. *Antisense Nucleic Acid Drug Dev.* 12:165-  
175, (2002).

Warren TL and Weiner GJ. Synergism between cytosine-guanine  
oligodeoxynucleotides and monoclonal antibody in the treatment of  
lymphoma. *Semin. Oncology*, 29:93-97 (2002).

20 Weeratna R et al., CpG DNA induces stronger immune responses with less  
toxicity than other adjuvants. *Vaccine*, 18, 1755-1762, (2000).

- Wernette CM et al., CpG oligodeoxynucleotides stimulate canine and feline immune cell proliferation. *Vet. Immunol. Immunopathol.* 84:223-236 (2002).
- Wooldridge JE et al., Immunostimulatory oligodeoxynucleotides containing CpG motifs enhance the efficacy of monoclonal antibody therapy of lymphoma. *Blood*, 89, 2994-2998, (1997).
- Yamamoto S et al., In vitro augmentation of natural killer cell activity and production of interferon alpha/beta and gamma with deoxyribonucleic acid fraction from *Mycobacterium bovis* BCG. *Jpn. J. Cancer Res.*, 79:866-873, (1988).
- 10 Yamamoto S. et al., DNA from Bacteria, but Not from Vertebrates, Induces Interferons, Activates Natural Killer Cells, and Inhibits Tumor Growth. *Microbiol. Immunol.* 36, 9:983-997, (1992).
- Yamamoto S. et al., Unique Palindromic Sequences in Synthetic Oligonucleotides are Required to Induce INF and Augment INF-Mediated Natural Killer Activity. *J. Immunol.*, 148, 12:4072-4076, (1992).
- 15 Yamamoto S., Mode of Action Of Oligonucleotide Fraction Extracted From *Mycobacterium bovis* BCG. *Kekkaku*, 69, 9:29-32, (1994).
- Yamamoto T et al., Ability of Oligonucleotides with Certain Palindromes to Induce Interferon Production and Augment Natural Killer Cell Activity Is Associated with Their Base Length. *Antisense Res. Devel.*, 4:119-123, 20 (1994).

- Yamamoto T et al., Lipofection of Synthetic Oligodeoxyribonucleotide Having a Palindromic Sequence AACGTT to Murine Splenocytes Enhances Interferon Production and Natural Killer Activity. *Microbiol. Immunol.* 38, 10:831-836, (1994).
- 5 Yamamoto T et al., Synthetic Oligonucleotides with Certain Palindromes Stimulate Interferon Production Of Human Peripheral B Lymphocytes in vitro, *Jpn. J. Cancer Res.*, 85:775-779, (1994).
- Yi A et al., IFN- $\gamma$  Promotes IL-6 and IgM Secretion in Response to CpG Motifs in Bacterial DNA and Oligonucleotides. *J. Immunol.* 558-564  
10 (1996).
- Yi AK et al. Role of mitogen-activated protein kinases in CpG DNA-mediated IL-10 and IL-12 production: central role of extracellular signal-regulated kinase in the negative feedback loop of the CpG DNA-mediated Th1 response. *J. Immunol.* 168:4711-4720, (2002).
- 15 Yi Ae-Kyung et al., Rapid Immune Activation by CpG Motifs in Bacterial DNA. *J. Immunol.*, 5394-5402 (1996).
- Yu D et al., Immunomers novel 3'-3'-linked CpG oligodeoxyribonucleotides as potent immunomodulatory agents. *Nucleic Acids Res.* 30:4460-4469. (2002).
- Zelenay, S. Et al., Immunomodulatory effects of plasmid DNA and synthetic  
20 oligodeoxynucleotides. *J. Eur. J. Immunol.* 33: 1382-92 (2003)
- Bylund, et al. *European Journal of Immunology* (2002), 32(10), 2847-2856
- Dalpke, et al. *Immunology* (2002), 106(1), 102-112

- Kandimalla, et al. Bioorganic & Medicinal Chemistry (2001), 9(3), 807-813
- Lang, et al. European Journal of Immunology (1999), 29(11), 3496-3506
- McCluskie, et al. Vaccine (2001), 19(17-19), 2657-2660
- Monteith, et al. Anti-Cancer Drug Design (1997), 12(5), 421-432
- 5 Oumouna, et al. Developmental & Comparative Immunology (2002), 26(3),  
257-269
- Parronchi, et al. Journal of Immunology (1999), 163(11), 5946-5953
- Vollmer, et al. Antisense & Nucleic Acid Drug Development (2002), 12(3),  
165-175
- 10 Zhao, et al. Biochemical Pharmacology (1996), 51(2), 173-82

## BACKGROUND OF THE INVENTION

### The immune system

The major function of the immune system is to protect the host from invading  
15 pathogens. A number of different cell types, both antigen-independent and  
antigen-specific, have evolved to detect and neutralize these invading  
pathogens. Among them, lymphocytes have an important characteristic,  
which is their ability to specifically recognize antigens, a feature not  
possessed by any other cell. This means that any lymphocyte function  
20 stimulated by an antigen is directed solely at that antigen.

Lymphocytes may be divided into two major populations: T and B. T-  
lymphocytes have a central role in regulating the immune response and for

this they produce and secrete lymphokines (i.e.interleukins). On the other hand, B-lymphocytes are the only cells that produce antibodies, which are proteins that recognize and bind antigens.

Some T-lymphocytes are known as helpers (Th-lymphocytes) because they  
5 assist B-cells to produce antibodies. T-lymphocytes express a characteristic membrane molecule designated CD4. Other T-lymphocytes are known as cytotoxic (CTL) because they are capable of killing certain cells. They express a different characteristic membrane protein designated CD8.

In mice, Th-lymphocytes have been subdivided in groups which are  
10 designated Th0, Th1 and Th2, according to the lymphokines they produce. In general, Th1 lymphocytes produce lymphokines which stimulate macrophages and CTLS (IL2, IFN  $\gamma$ , TNF- $\beta$ ); Th2 lymphocytes produce lymphokines which stimulate B-lymphocytes to proliferate and to produce antibodies (IL 2, IL5, IL6, IL10, IL13); and Th0 lymphocytes produce a mixture  
15 of lymphokines and are thought to be an intermediate stage from which Th1 and Th2 lymphocytes are derived. In humans, Th1- and Th2- like lymphocytes have been demonstrated, although they seem to show a less strict division with respect to their patterns of cytokine secretion.

A third population of lymphocytes, which lack the major makers of T and B  
20 cells include the natural killer cells (NK cells), the killer cells (K cells) and the lymphokine-activated killer cells (L A K cells). NK cells can kill certain tumor cells and some virally infected cells, but unlike cytotoxic T-lymphocytes, they

are not capable of recognizing a specific antigen. First, they can bore a hole in a target cell by secreting perforin molecules to form a membrane attack complex on the surface of the target. NK cells can then secrete enzymes that enter the target cell and cause it to commit suicide. As a second mechanism of destruction, Fas ligand, a protein present on NK cells can interact with Fas, another protein present on the surface of the target cell, inducing the target cell to commit suicide by apoptosis. NK cells are also able to bind to cells, which have antibody to them via their antigen-binding regions and kill them. L A K cells do not specifically recognize an antigen but they are capable of destroying a wider range of targets than NK cells.

Macrophages and dendritic cells play a critical role in initiating immune responses, helping T cells to respond to antigens.

There are several classes of antibody. The IgG class comprises most of the circulating antibodies and it has four subclasses designated IgG1, IgG2, IgG3 and IgG4. The IgM class comprises about 10 % of the circulating antibodies. These are the principal antibodies produced during the primary immunological response. The IgA class comprises most of the antibodies secreted at mucous membranes and exerts its protective effect by blocking access of the antigen to the inner body. The IgD class comprises less than 1 % of serum antibodies and its biological role is largely unknown. The IgE class comprises antibodies that are mainly bound to the surface of most cells



and basophils. These antibodies are associated with reactions that occur in individuals who are undergoing allergic reactions.

#### Vaccines and vaccine adjuvants

- 5 Vaccines are preparations used to stimulate animals to mount an immune response against antigens included in the vaccine.

Vaccines often include adjuvants, which are substances that used in combination with specific antigen produce more immunity than the antigen used alone. (Ramon, G., 1926. Procédes pour accroître la production des  
10 antitoxins. Ann. Inst. Pasteur. 40, 1- 10).

Many kinds of compounds function as vaccine adjuvants (Edelman, R., 2000. An overview of adjuvant use, in : Vaccine Adjuvants. Preparation Methods and Research Protocols. D.T. O' Hagan, Ed., Humana Press, Totowa, New Jersey. References cited in this article are incorporated herein as background  
15 material).

However, currently, the only adjuvants approved for use in humans are aluminum salts (Gupta, R.K. and Rost, B.E., 2000. Aluminum compounds as vaccine adjuvants in: Vaccine Adjuvants. Preparation Methods and Research Protocols. D.T. O' Hagan, Ed., Humana Press, Totowa, New Jersey) and the  
20 oil-in-water emulsion M F 59 (Ott, G. Radhakrishnan, R. Fang, J. and Hora, M., 2000. The adjuvant M F 59: A 10- Year Perspective, in : Vaccine

Adjuvants. Preparation Methods and Research Protocols. D.T. O' Hagan, Ed., Humana Press, Totowa, New Jersey).

Nucleic acids as immunostimulatory compounds

5 Several polynucleotides have been demonstrated to have immunostimulatory properties. For example, poly (I,C) is an inducer of interferon (IFN) production, macrophage activation and NK cell activation (Talmadge, J.E., Adams, J., Phillips, H., Collins, M., Lenz, B., Schneider, M., Schlick, E., Ruffmann, R., Wilttrout, R.H., Chirigos, M.A.1985. Immunomodulatory effects  
10 in mice of polyinosinic-polycytidylic acid complexed with poly-L:-lysine and carboxymethylcellulose. Cancer Res. 45:1058; Wilttrout, R.H., Salup, R.R., Twilley, T.A., Talmage, J.E. 1985. Immunomodulation of natural killer activity by polyribonucleotides. J. Biol. Resp. Mod. 4:512), poly (dG,dC) is mitogenic for B cells (Messina, J.P., Gilkerson, G.S., Pisetsky, D.S. 1993. The influence  
15 of DNA structure on the in vitro stimulation of murine lymphocytes by natural and synthetic polynucleotide antigens. Cell. Immunol. 147:148) and induces IFN and NK activity (Tocunaga, T., Yamamoto, S., Namba, K.1988. A synthetic single-stranded DNA, poly(dG,dC), induces interferon- $\alpha/\beta$  and - $\gamma$ , augments natural killer activity, and suppresses tumor growth. Jpn. J. Cancer  
20 Res. 79:682).

Bacterial DNA has also been reported to have immunostimulatory properties. These properties include the induction of cytokines (interferon gamma (IFN  $\gamma$ ),

alpha (IFN  $\alpha$  ) and beta (IFN  $\beta$ )), tumor necrosis factor alpha (TNF  $\alpha$  ), interleukin 6 (IL6), 12 (IL 12) and 18 (IL 18), as well as the direct stimulation of B cells (Yamamoto, S. et al. 1988. In vitro augmentation of natural killer cell activity of interferon  $\alpha/\beta$  and  $\gamma$  with deoxyribonucleic acid fraction from *Mycobacterium bovis* BCG. Jpn. J. Cancer Res. (Gann) 79: 866-873; Yamamoto S. et al., 1992. DNA from bacteria, but not from vertebrates, induces interferons, activates natural killer cells and inhibits tumor growth. Microbiol. Immunol. 36: 983-997; Klinman, D.M., Yi, A-K., Beaucage, S.L., Conover, J. and Krieg, A.M.,1996. CpG motifs present in bacterial DNA rapidly induce lymphocytes to secrete interleukin 6, interleukin 12 and interferon  $\gamma$ . Proc. Natl. Acad. Sci. USA 93, 2879-2883. Halpern, M. D., et al. 1996. Bacterial DNA induces murine interferon - $\gamma$  production by stimulation of interleukin -12 and tumor necrosis factor - $\alpha$ . Cell. Immunol. 167: 72-78. Sparwasser, T. et al.,1997. Macrophages sense pathogens via DNA motifs: induction of tumor necrosis factor - $\alpha$  - mediated shock. Eur. J. Immunol. 27: 1671-1679; Krieg, A.M. et al., 1995. CpG motifs in bacterial DNA trigger direct B-cell activation. Nature 374: 345-349).

In contrast, it has been reported that mammalian DNA has no significant immune effects (Pisetsky, D.S. 1996. The immunologic properties of DNA. J. Immunol. 156: 421-423; Messina et al. 1991. Stimulation of in vitro murine lymphocyte proliferation by bacterial DNA. J. Immunol. 147: 1759).

Synthetic DNA has also been reported to be immunostimulatory if it contains unmethylated CpG motifs. (Yamamoto, S et al.; 1992. Unique palindromic sequences in synthetic oligonucleotides are required to induce INF and augment INF-mediated natural killer activity. J. Immunol. 148: 4072-4076;

5 Ballas, Z.K., et al.; 1996. Induction of NK activity in murine and human cells by CpG motifs in oligodeoxynucleotides and bacterial DNA. J. Immunol. 157: 1840-1845; Hartmann, G., Krieg, A.M. 2000. Mechanism and function of a newly identified CpG DNA motif in human primary B cells. J. Immunol. 164:944; Hartmann, G., Weeratna, R.D., Ballas, Z.K., Payette, P., Blackwell,


10 S., Suparto, I., Rasmussen, W.L., Waldschmidt, M., Sajuthi, D., Purcell, R.H., Davis, H.L., Krieg, A.M. 2000. Delineation of a CpG phosphorothioate oligodeoxynucleotide for activating primate immune responses in vitro and in vivo. J. Immunol. 164:1617; Verthelyi, D., Ishii, K.J., Gursel, M., Takeshita, F., Klinman, D.M. 2001. Human peripheral blood cells differentially recognize and

15 respond to two distinct CpG motifs. J. Immunol. 166:2372). However, one oligonucleotide containing phosphorothioate bonds that lack CpG motifs has been found to have some immunostimulatory activity on human B cells (Liang, H., Nishioka, Y., Reich, C.F., Pisetsky, D.S., Lipsky, P.E. 1996. Activation of human B cells by phosphorothioate oligonucleotides. J. Clin.

20 Invest. 98:1119). This particular non-CpG oligonucleotide containing phosphorothioate bonds is a poly-T chain, 20 nucleotides long. Also, Vollmer et al (Vollmer J, Janosch A, Laucht M, Ballas ZK, Schetter C, Krieg AM.

Highly immunostimulatory CpG-free oligodeoxynucleotides for activation of human leukocytes. *Antisense Nucleic Acid Drug Dev.* 12:165-175, 2002) reported immunostimulation by phosphorothioate poly-T ODNs. These authors pointed out that poly-T ODNs are only active as phosphorothioate ODNs and have much lower activity than CpG ODNs.

It has now been discovered that non-CpG oligonucleotides containing the following non-palindromic sequence motif:



wherein  $X_1$  is C or T,  $X_2$  is C, T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein C never precedes G (in other terms, the nucleic acid motif does not consist of a CpG oligonucleotides) have potent immunostimulatory activity in animals of the order *Primate* including humans. Therefore, these oligonucleotides can be administered to subjects to treat "immune system deficiencies" or used as adjuvants, in conjunction with a vaccine, to boost the immune system in order to have a better response to the vaccine. They can also be administered to subjects to increase their responsiveness to tumors.

#### SUMMARY OF THE INVENTION

It has now been discovered that non-CpG oligonucleotides containing the following non-palindromic sequence motif:



wherein  $X_1$  is C or T,  $X_2$  is C, T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are Ts and wherein C never precedes G (in other terms, the nucleic

acid motif does not consist of a CpG oligonucleotides), have the ability to stimulate the immune response of animals of the order *Primate*, including humans.

According to a preferred embodiment,  $X_1$  consists of a C. It is preferable if  
5  $X_3, X_4, X_5, X_6, X_7, X_8$  of the immunostimulatory motif consists of TTTTGT, ATTTGT or ATTGGT. It is even more advantageous if  $X_1$  is C;  $X_3$  is T or A;  $X_4$  is T;  $X_5$  is T;  $X_6$  is T or G;  $X_7$  is G and  $X_8$  is T.

The oligonucleotides of this invention are useful as adjuvants in a vaccine formulation comprising one or more antigens. In embodiments of this aspect,  
10 the vaccine formulation can be liquid or lyophilized in dosage form. Many dosage forms are known in the art and can be applied herein. In embodiments of this aspect the oligonucleotides of this invention are present in the composition at a dose of from about 10 to 10,000  $\mu\text{g}$  per dose. In these preparations, the oligonucleotides of this invention may be combined with  
15 other immunostimulant compounds. Examples of well-known immunostimulants are:  $\alpha$ -interferon,  $\beta$ -interferon,  $\gamma$ -interferon, granulocyte macrophage colony stimulator factor (GM-CSF), interleukin 2 (IL2), interleukin 12 (IL12) and CpG oligonucleotide.

In preferred embodiments, the antigenic component of the vaccine is one or  
20 more antigens, either natural or recombinant, of viruses like: Human immunodeficiency viruses, such as HIV-1 and HIV-2, polio viruses, hepatitis A virus, human coxsackie viruses, rhinoviruses, echoviruses, equine encephalitis viruses, rubella viruses, dengue viruses, encephalitis viruses,

yellow fever viruses, coronaviruses, vesicular stomatitis viruses, rabies viruses, ebola viruses, parainfluenza viruses, mumps virus, measles virus, respiratory syncytial virus, influenza viruses, Hantaan viruses, bunga viruses, hemorrhagic fever viruses, reoviruses, orbiviruses, rotaviruses, Hepatitis B virus, parvoviruses, papilloma viruses, polyoma viruses, adenoviruses, herpes simplex virus (HSV) 1 and 2, varicella zoster virus, cytomegalovirus (CMV), variola viruses, vaccinia viruses, pox viruses, African swine fever virus, the unclassified agent of delta hepatitis, the agents of non-A, non-B hepatitis; or one or more antigens of infectious bacteria like: *Helicobacter pylori*, *Borrelia burgdorferi*, *Legionella pneumophila*, *Mycobacterium tuberculosis*, *Mycobacterium bovis* (BCG), *Mycobacterium avium*, *Mycobacterium intracellulare*, *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Listeria monocytogenes*, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Klebsiella pneumoniae*, *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Clostridium perfringens*, *Clostridium tetani*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pasteurella multocida*, and *Treponema pallidum*; of infectious fungi like: *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis*, *Candida albicans*; of infectious protists like: *Plasmodium falciparum*, *Trypanosoma cruzi*, *Leishmania donovani* and *Toxoplasma gondii*. and; of human tumoral cells.

In embodiments of this aspect, one or more of the oligonucleotides of this invention and the antigen are administered simultaneously locally (by oral, rectal, intranasal or transdermal route) or systemically (by intradermic or intramuscular injection).

- 5 An aspect of this invention is the use of the present oligonucleotides for the manufacture of a medicament for vaccinating a person. The person can be vaccinated *prophylactically* or *therapeutically*.

A *prophylactic* vaccine is designed to elicit protection from a disease caused by an infectious agent through the induction of specific immunity.

- 10 A *therapeutic* vaccine is designed to induce remission of an illness (i.e. a tumor and metastasis or illness associated with an infectious agent like the human immunodeficiency virus).

The method of vaccination includes administering one or more of the oligonucleotides of this invention and one or more antigens – that is, the vaccine can be designed against one disease target or a combination of  
15 disease targets.

Another aspect of this invention is a method of treatment of a person with a tumoral disease or an immunological disorder, namely the use of one or more of the oligonucleotides of this invention for the manufacture of a medicament  
20 to stimulate his/her endogenous immune response. Examples of tumoral disease are: Chronic Myelogenous Leukemia, Precursor B-lymphoblastic lymphoma, B-cell chronic lymphocytic leukaemia, Lymphoplasmacytic



lymphoma, Mantle cell lymphoma, Follicle center lymphoma, (follicular and diffuse), Marginal zone-B lymphoma, Extranodal lymphoma, Nodal marginal zone B-cell lymphoma, Splenic marginal zone B-cell lymphoma, Hairy cell leukaemia, Plasmacytoma, Diffuse large B-cell lymphoma, Burkitt's lymphoma, High grade B-cell lymphoma, Burkitt like, Melanoma, Kaposi's Sarcoma, Multiple Myeloma, Renal Cell Carcinoma, Bladder Cancer, Lung Cancer, Skin Cancer, Breast Cancer, Colon Cancer and Uterus Cancer. Examples of immunological disorders are: Allergy, Severe Combined Immunodeficiency, Chronic Granulomatous disease, and Acquired Immunodeficiency Disease.

In embodiments of this aspect, one or more of the oligonucleotides of this invention are present in a pharmaceutical formulation that can be liquid or lyophilized in dosage form. Many dosage forms are known in the art and can be applied herein. In embodiments of this aspect one or more of the oligonucleotides of this invention are present in the composition at a dose of from about 10 to 10,000  $\mu\text{g}$  per dose. In these preparations one or more of the oligonucleotides of this invention may be combined with other immunostimulant compounds. Examples of well-known immunostimulants are:  $\alpha$ -interferon,  $\beta$ -interferon,  $\gamma$ -interferon, granulocyte macrophage colony stimulator factor (GM-CSF), interleukin 2 (IL2), interleukin 12 (IL12), CpG oligonucleotides and *Mycobacterium bovis* BCG cells. Also, one or more of the oligonucleotides of this invention may be combined with an antiinfective or

anticancer drug, or a surgical procedure. In all these cases, the oligonucleotides of this invention may be administered before, after or simultaneously with the alternative treatment.

Another aspect of this invention is that a person with a tumoral disease or an immunological disorder could be treated by contacting lymphocytes or plasmacytoid dendritic cells from the subject, with one or more of the oligonucleotides of this invention "ex vivo" and readministering the activated cells to the subject. In embodiments of this aspect one or more of the oligonucleotides of this invention are present in the incubation media in a concentration of about 0.10 to 100 µg per ml.

#### BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 is a graph plotting the proliferation index of human peripheral blood mononuclear leukocytes (PBMC) incubated with the phosphorothioate ODN IMT 023 (Seq. ID N° 9) or the phosphorothioate CpG ODN 2006:

5' TCGTCGTTTTGTCGTTTTGTCGTT 3' or the phosphorothioate non-CpG ODN IMT 021 (SEQ ID NO:1). Data represent the mean and standard deviation of five independent assays.

Fig 2 shows the results from a flow-cytometric study using human peripheral blood mononuclear leukocytes (PBMC) to determine the comparative effect of the phosphorothioate CpG ODN 2006:

5' TCGTCGTTTTGTCGTTTTGTCGTT 3'

or the phosphorothioate non-CpG ODN IMT 021 (SEQ ID NO:1) on activation of the B cell population. B cell activation was measured by the induction of CD86 on CD69 positive cells.

(a) Representative flow-cytometric diagrams comparing B cell activation by ODNs IMT023 (Seq. ID N<sup>o</sup> 9), 2006 or IMT021 (SEQ ID NO:1); b) Representative histograms of the flow-cytometric diagrams showed in (a).

Fig. 3 is a graph plotting the proliferation index of PBMC incubated with the indicated phosphorothioate ODNs. Data represent the mean and standard deviation of five independent assays.

Fig. 4. shows the induction of CD86 in CD19+ (B) cells and CD69 in CD56 + (NK) cells by non-CpG ODNs claimed in WO 01/22972 . Human PMBC were cultured for 48 hr with indicated *phosphodiester* ODNs or controls and then stained with fluorescent anti-CD19/anti-CD86 (a) or, anti-CD19/anti-CD40 (b) or, anti-CD19/anti-MHC I (c). Flow cytometric results are presented as histograms corresponding to cells in the double positive gate. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (O): phosphodiester ODN. ODN 2080, a CpG ODN of the following sequence: 5'-TCGTCGTTCCCCCCCCCCCC-3' was used as positive control.

Fig. 5 shows the influence of the position of the immunostimulatory sequence motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed on the immunostimulatory activity of

non-CpG ODNs measured in PMBC proliferation assays. Data represent the mean and standard deviation of three independent assays performed in quadruplicate.

Fig. 6 shows the induction of CD40, MHC I and MHC II in CD19+ cells (B cells) by non-CpG-ODNs bearing the immunostimulatory sequence motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. Human PMBC were cultured for 24 hr with indicated ODNs and then stained with fluorescent anti-CD19/anti-CD40 (a) or, anti-CD19/ anti-MHC I (b) or, anti-CD19/ anti-MHC II (c). Flow-cytometric results are presented as histograms corresponding to CD19+ cells. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (S) means phosphorothioate ODN.

Fig. 7 shows the induction of CD86, CD40 and MHC I in purified B cells by non-CpG-ODNs bearing the immunostimulatory sequence motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. Human purified B cells were cultured for 24 hr with indicated ODNs and then stained with fluorescent anti-CD19/anti-CD86 (a) or, anti-CD19/anti-CD40 (b) or, anti-CD19/anti-MHC I (c). Flow-cytometric results are presented as histograms corresponding to CD19+ cells. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (S) means phosphorothioate ODN.

Fig. 8 shows induction of CD86, CD40 and MHC I in CD19+ cells (B cells) by phosphodiester non-CpG-ODNs bearing the immunostimulatory sequence motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. Human PMBC were cultured for 48 hr with indicated phosphodiester ODNs and then stained with fluorescent anti-CD19/anti-CD86 (a) or, anti-CD19/anti-CD40 (b) or, anti-CD19/anti-MHC I (c). Flow-cytometric results are presented as histograms corresponding to CD19+ cells. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (O) means phosphodiester ODN.

Fig. 9 shows stimulation of PMBC proliferation by non-CpG ODNs bearing the motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed in non-human primates. *Cebus apella* or *Macacca fascicularis* PMBC were cultured for 72 hr with indicated ODNs (6µg/ml). Data represent the mean and standard deviation of four replicates.

Fig. 10 shows the induction of CD86, CD40 and MHC I in CD 19+ (B cells) of a patient suffering B cell chronic lymphocytic leukemia (CLL) by non-CpG-ODNs bearing the motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. PMBC were cultured for 24hr with indicated ODNs and then stained with fluorescent anti-CD19/anti-CD86 (a) or, anti-CD19/anti-CD40 (b) or, anti-CD19/anti-MHC I (c). Flow-cytometric results are presented as histograms corresponding to CD19+ cells. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (S) means phosphorothioate ODN.

Fig. 11 shows the stimulation of malignant B cell apoptosis by non-CpG-ODNs bearing the motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. PMBC were cultured for 14 hr. or 5 days with indicated ODNs and then stained with propidium iodide and annexin V-FITC using the ANNEXIN V:FITC assay kit from Serotec (Raleigh, NC, USA). Flow-cytometric results are presented as a dot-plot of annexin V-FITC fluorescence versus propidium iodide (PI) fluorescence. This plot show three cell populations: a) low PI-low annexin that are viable cells, b) low PI-high annexin that are apoptotic cells and, c) high PI-high annexin that are secondary necrotic cells.

Fig. 12 shows the induction of CD86, CD40 and MHC I in purified plasmacytoid dendritic cells by non-CpG-ODNs bearing the motif  $X_1X_2X_3X_4X_5X_6X_7X_8$  here disclosed. More than 95 % pure plasmacytoid dendritic cells were cultured for 24hr with indicated ODNs and then stained with fluorescent anti-CD4/anti-CD11c/anti-CD86 (a) or, anti-CD4/anti-CD11c /anti-CD40 (b) or, anti-CD4/anti-CD11c /anti-MHC I (c). Flow cytometric results are presented as histograms corresponding to CD4+CD11c- cells. Open histograms correspond to cells cultured in absence of ODN and shaded histograms to cells cultured in presence of ODN. ODN (S) means phosphorothioate ODN.

Fig. 13 is a graph plotting the IL-10 levels in supernatants of five independent human peripheral blood mononuclear leukocytes (PBMC) cultured 72 hr with

the phosphorothioate non-CpG ODN IMT 504 (Seq. ID N<sup>o</sup> 2.), the phosphorothioate CpG ODN 2006: 5' TCGTCGTTTTGTCGTTTTGTCGTT 3', or the phosphorothioate non-CpG ODN IMT 022 (SEQ ID NO: 175). Data represent the mean and standard deviation of five independent assays.

- 5 Fig. 14 is a graph plotting the IL-5 levels in supernatants of five independent human peripheral blood mononuclear leukocytes (PBMC) cultured 96 hs with the phosphorothioate non-CpG ODN IMT 504 (Seq. ID N<sup>o</sup> 2.), the phosphorothioate CpG ODN 2006: 5' TCGTCGTTTTGTCGTTTTGTCGTT 3', or the phosphorothioate non-CpG ODN IMT 022 (SEQ ID NO: 175). Data  
10 represent the mean and standard deviation of five independent assays.

#### DETAILED DESCRIPTION OF THE INVENTION

##### Definitions:

- An "allergy" refers to acquired hypersensitivity to a substance (allergen).  
15 Examples of allergies are eczema, allergic rhinitis, asthma and urticaria.
- An "immune system deficiency" refers to a disease in which the immune system is not functioning in normal capacity.
- As used herein, the term "oligonucleotide" or "oligo" shall mean multiple nucleotides (i.e. molecules comprising a sugar (e.g. ribose or deoxyribose)  
20 linked to a phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g. cytosine (C), thymine (T) or uracil (U)) or a substituted purine (e.g. adenine (A) or guanine (G)). The term

"oligonucleotide" as used herein refers to both oligoribonucleotides (ORNs) and oligodeoxyribonucleotides (ODNs). The term "oligonucleotide" shall also include oligonucleosides (i.e. an oligonucleotide minus the phosphate) and any other organic base containing polymer. Oligonucleotides can be obtained from existing nucleic acid sources (e.g. genomic or cDNA), but are preferably synthetic (e.g. produced by oligonucleotide synthesis).

- An "oligonucleotide" refers to multiple nucleotides linked by phosphodiester bonds.

- An "immunostimulatory oligonucleotide" refers to an oligonucleotide which stimulates (i.e. has a mitogenic effect on, or induces or increases or decreases cytokine expression by) a cell of the immune system (i.e.: a lymphocyte, a macrophage) in a statistically significant manner.

- A "Strong immunostimulatory oligonucleotide" refers to an oligonucleotide which stimulates "in vitro" cells of the immune system of an animal of the order Primate with an activity of at least 60% of the immunostimulatory activity of the well known CpG ODN 2006, under the same experimental conditions and using identical chemical backbone (i.e. phosphorotioate backbone or phosphodiester backbone).

- A "CpG" refers to a cytosine-guanine dinucleotide.

- A "CpG oligonucleotide" refers to an oligonucleotide which stimulates a cell of the immune system and its immunostimulatory activity critically depends on the presence of at least one CpG in its sequence.



- A "non-CpG oligonucleotide" refers to an oligonucleotide which stimulates a cell of the immune system and its immunostimulatory activity does not critically depends on the presence of a CpG in its sequence.
- A "subject" refers to an animal of the order Primate, including humans.
- 5 - As used herein, the term "treating" refers to a process by which the symptoms of a disease, and more particularly, infectious diseases, tumoral diseases or immunological disorders are ameliorated or completely eliminated.
- As used herein, the term "preventing" refers to a process by which a  
10 disease, and more particularly infectious diseases or tumoral diseases or immunological disorders are obstructed or delayed.
- In a preferred embodiment, the immunostimulatory oligonucleotides of the invention are advantageously modified into stabilized oligonucleotides. Such stabilized immunostimulatory oligonucleotide may be particularly useful to  
15 obtain a prolonged immunostimulation. As used herein, a "stabilized oligonucleotide" refers to an oligonucleotide that is relatively resistant to in vivo degradation (e.g. via an exo- or endo-nuclease). Preferred stabilized oligonucleotides of the present invention comprise a phosphate backbone modification. More particularly, the phosphate backbone modification is  
20 preferably a 5' inter-nucleotide linkage modification, for instance, at the first two nucleotides of the 5' end of the oligonucleotide of the invention. Furthermore, the phosphate backbone modification may be a 3' inter-

nucleotide linkage modification. In such a case, the modification may occur, for instance, at the last two nucleotides of the 3' end of the oligonucleotide of the invention. Even more preferably, the immunostimulatory oligonucleotide of the invention may be stably modified so as to comprise a  
5 phosphorothioate-linked nucleotide (i.e. at least one of the phosphate oxygens is replaced by sulfur). In the most preferred embodiment, most if not all the nucleotides of the immunostimulatory oligonucleotides of the invention comprise a phosphorothioate-linked nucleotide.

Other stabilized oligonucleotides may alternatively include: nonionic DNA  
10 analogs, such as alkyl- and aryl- phosphonates (in which the charged phosphonate oxygen is replaced by an alkyl or aryl group), phosphodiester and alkylphosphotriesters, in which the charged oxygen moiety is alkylated. Oligonucleotides which contain a diol, such as tetraethyleneglycol or hexaethyleneglycol, at either or both termini have also been shown to be  
15 substantially resistant to nuclease degradation.

The present invention provides methods to augment the immune response of animals of the order *Primate*, including humans, adding one or more of the oligonucleotides of this invention to vaccines, or performing a treatment based on the administration of one or more of the oligonucleotides of this  
20 invention to a person with a tumoral disease or an immunological disorder, or contacting "ex vivo" white blood cells obtained from a person with a tumoral disease or an immunological disorder with one or more of the

oligonucleotides of this invention and readministering these activated white blood cells to the same person.

Vaccine compositions containing one or more of the oligonucleotides of this invention can present antigens directly (i.e. in the form of a defined protein or polysaccharide) or as a part of a complex biological entity (i.e. complete  
5 viruses; complete bacterial cells; bacterial membranes or artificial conjugates like polysaccharide-protein conjugates). These antigens can be combined in multiple vaccines.

A vaccine composition including at least one antigen is formulated to include  
10 one or more of the oligonucleotides of this invention.

For example the antigen can be *Moraxella catharralis* killed cells or subcellular fractions of these cells or *Hepatitis B virus* surface antigen, either natural or produced by means of the DNA recombinant technology.

One or more of the oligonucleotides of this invention may be formulated alone  
15 or together with one or more antigens in a pharmaceutical composition, which may also include carriers, thickeners, diluents, buffers, preservatives, surface active agents, anti-microbial agents, anti-inflammatory agents, anesthetics and the like.

The formulation can be liquid or lyophilized. The pharmaceutical composition  
20 may be administered in a number of ways depending of whether local or systemic treatment is desired, and on the area to be treated. Administration may be done topically, orally, by inhalation or parenterally. Formulation for

topical administration may include ointments, lotions, creams, gels; drops, suppositories, sprays, liquids and powders. Pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, tablets and the like. Thickeners, flavorings, diluents, emulsifiers and the like may be necessary or desirable. Formulations for parenteral administration include sterile aqueous solutions, which may also contain buffers, diluents and other additives. A vaccine containing one or more antigens and one or more of the oligonucleotides of this invention can be formulated and used for prophylactic or therapeutic purposes.

Common antigens used in viral prophylactic vaccines are from *Hepatitis B virus*, *Hepatitis A virus* and *Influenza virus*. Common antigens used in bacterial prophylactic vaccines are from *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catharralis*, *Klebsiella pneumoniae* and *Mycobacterium bovis* (BCG). Common antigens used in therapeutic vaccines are from *Papilloma virus*, *HIV virus* and melanoma cells.

A further refinement of a vaccine formulation is to incorporate one or more of the oligonucleotides of this invention as adjuvant/s and the antigen/s into a delivery vehicle to provide for delayed release of the active compounds of the vaccine over time. This can be accomplished by various means known in the art. Examples of these means are encapsulation into Poly (lactide-coglicolide)

- micro particles (Kersten, G.F.A. and Gander, B. 1996. Biodegradable Micro Spheres as vehicles for antigens, in: S.H.E. Kaufmann, ed. Concepts in Vaccine Development. Walter de Gruyter. Berlin-New York), liposomes (Gregoriadis, G. et al. 2000. Liposomes as Immunological Adjuvants and Vaccine Carriers, in: S.H.E. Kaufmann, ed. Concepts in Vaccine Development. Walter de Gruyter. Berlin-New York) and poly (methyl methacrylate) nanoparticles (Kreuter, J. 2000. Poly (Methyl Methacrylate) nanoparticles as vaccine adjuvants, in: S.H.E. Kaufmann, ed. Concepts in Vaccine Development. Walter de Gruyter. Berlin-New York).
- 10 Another refinement of the vaccine formulation is to conjugate the antigen/s and one or more of the oligonucleotides of this invention, by chemical means (Mier W, Eritja R, Mohammed A, Haberkorn U, Eisenhut M.2000. Preparation and evaluation of tumor-targeting peptide-oligonucleotide conjugates. Bioconjug. Chem. 11:855).
- 15 Many vaccine formulations are known in the art and can be used by substituting one or more of the oligonucleotides of this invention for the adjuvant previously known or by simply adding one or more of the oligonucleotides of this invention to the original formulation.
- Based on their immunostimulatory properties, one or more of the
- 20 oligonucleotides of this invention can also be administered to a subject in vivo to treat a tumoral disease or an immune system disorder.

Examples of common tumoral diseases are: Chronic Myelogenous Leukemia, Melanoma, Kaposi's Sarcoma, Multiple Myeloma, Renal Cell Carcinoma, Bladder Cancer, Lung Cancer, Skin Cancer, Breast Cancer, Colon Cancer and Uterus Cancer. Examples of common immunological disorders are:

5 Allergy, Severe Combined Immunodeficiency, Chronic Granulomatous disease, and Acquired Immunodeficiency Disease.

The pharmaceutical composition for these treatments may include one or more of the oligonucleotides of this invention together with carriers, thickeners, diluents, buffers, preservatives, surface active agents, anti-  
10 microbial agents, anti-inflammatory agents, anesthetics and the like. The formulation can be liquid or lyophilized.

The pharmaceutical composition may be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Administration may be done topically, orally, by inhalation or  
15 parenterally.

Formulation for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Formulations for oral administration include powders or granules, suspensions or solutions in water  
20 or non-aqueous media, capsules, tablets and the like. Thickeners, flavorings, diluents, emulsifiers and the like may be necessary or desirable. Formulations for parenteral administration include sterile aqueous solutions, which may

also contain buffers, diluents and other additives. Alternatively, one or more of the oligonucleotides of this invention can be contacted "ex vivo" with immunocompetent cells (i.e. B cells or plasmacytoid dendritic cells) obtained from a subject having a tumoral disease or an immune system deficiency and activated cells can then be reintroduced in the subject.

As it will be apparent from the following examples the preferred oligonucleotides are those which stimulate "in vitro" cells of the immune system of an animal of the order Primate with an activity of at least 60% of the immunostimulatory activity of the well known CpG ODN 2006, under the same experimental conditions and using identical chemical backbone (i.e the strong immunostimulatory oligonucleotides).

Among these are preferred those having about 14 to 40 nucleotides and wherein  $X_3X_4X_5X_6X_7X_8$  consists of TTTTGT, the most preferred strong immunostimulatory oligonucleotide being IMT 504, namely TCATCATTTTGTTCATTTTGTTCATT (SEQ ID N°2).

#### EXAMPLE 1

##### Materials and Methods

The following materials and methods were generally used throughout the examples.

##### 1) Oligonucleotides

Oligonucleotides having phosphorothioate internucleotide linkages were purchased, purified by high-pressure liquid chromatography (HPLC), from Operon Technologies (Alameda, California) or Annovis (Aston, Pennsylvania) or Oligos Etc (Bethel, Maine). ODNs were suspended in depyrogenated  
5 water, assayed for LPS contamination using the Limulus test and kept at -20 °C until used. Purity was assessed by HPLC and PAGE assays. ODN preparations were used if LPS levels were undetectable.

## 2) Antibodies

Antibodies used in assays were purchased from Serotec (Raleigh, NC).

## 10 3) Peripheral blood mononuclear leukocytes (PBMC)

Blood was obtained by venipuncture from healthy donors using heparin as anticoagulant. PMBC were isolated by Ficoll-Hypaque (Sigma Diagnostics Inc., St. Louis, MO) density gradient centrifugation. Briefly, blood samples diluted 1:2 in RPMI-1640 medium (PAA laboratories GmbH, Linz, Austria)  
15 supplemented with 2.0 mM L-glutamine and 50.0 µg/ml gentamicin and 20 mM HEPES were centrifuged at 1000 × g for 40 minutes at 20 °C. PMBC were isolated, washed and suspended in medium supplemented with 10 % fetal calf serum.

## 4) Purification of cells

20 B lymphocytes and plasmacytoid dendritic cells were purified from human PMBC by positive selection using the MACS magnetic cell separation systems (Miltenyi Biotec, Germany).



### 5) Cell Proliferation assays

Blood was obtained by venipuncture from healthy donors using heparin as anticoagulant. PMBC were isolated by Ficoll-Hypaque (Sigma Diagnostics Inc., St. Louis, MO) density gradient centrifugation. Briefly, blood samples  
5 diluted 1:2 in RPMI-1640 medium (PAA laboratories GmbH, Linz, Austria) supplemented with 2.0 mM L-glutamine and 50.0 µg/ml gentamicin and 20 mM HEPES were centrifuged at  $1000 \times g$  for 40 minutes at 20 °C. PMBC were isolated, washed and suspended in medium supplemented with 10 % fetal calf serum.

10 PMBC were cultured in RPMI-1640 media supplemented with 10% (v/v) heat inactivated fetal calf serum (FCS), 2.0 mM L-glutamine and 50.0 µg/ml gentamicin.  $1 \times 10^5$  cells/well were incubated in 96 well microtiter plates (NUNC, Denmark) at 37 °C in a 5% CO<sub>2</sub> humidified atmosphere for 72 hr. PMBC were stimulated with ODNs at 0.375 µg/ml unless otherwise stated.  
15 Eighteen hours prior to cell harvest, 1 µCi of <sup>3</sup>H-Thymidine (Amersham, specific activity: 25 Ci/mmol) was added to each well. Cells were harvested onto glass-fiber filters and <sup>3</sup>H incorporation measured by scintillation counting. Standard deviations of quadruplicate wells were <10%.

### 6) IL6 Assay

20 PMBC ( $3 \times 10^5$ /well) were cultured as described above with ODNs (6 µg/ml) for 24 hr. After this, supernatants were collected and IL6 levels measured by

ELISA. Briefly, 96 well micro titer plates (NUNC, Denmark) were coated with anti-IL6 antibodies and blocked with RPMI 1640 media supplemented with 10% (v/v) heat inactivated FCS. IL6 was detected colorimetrically using biotin-labeled antibodies followed by peroxidase-conjugated strepto-avidin and then peroxidase-specific colorimetric substrate. Standard curves were generated using known amounts of recombinant IL6. The detection limit of these assays was 30 pg/ml. All assays were performed in duplicate.

#### 7) IL-10 Assay

PBMC ( $3 \times 10^5$ /well) were cultured as described above with ODNs (1.5  $\mu$ g/ml) for 72 hr. After this, supernatants were collected and IL-10 levels measured by ELISA. Briefly, 96 well micro titer plates (NUNC, Denmark) were coated with anti-IL-10 antibodies and blocked with RPMI 1640 media supplemented with 10% (v/v) heat inactivated FCS. IL-10 was detected colorimetrically using biotin-labeled antibodies followed by peroxidase-conjugated strepto-avidin and then peroxidase-specific colorimetric substrate. Standard curves were generated using known amounts of recombinant IL-10. The detection limit of these assays was 20 pg/ml. All assays were performed in duplicate.

#### 8) IL-5 Assay

PBMC ( $3 \times 10^5$ /well) were cultured as described above with ODNs (1.5  $\mu$ g/ml) and IL-4 (5 ng/ml) for 72 hr. After this, supernatants were collected and IL-5 levels measured by ELISA. Briefly, 96 well micro titer plates (NUNC, Denmark) were coated with anti-IL-5 antibodies and blocked with RPMI 1640

media supplemented with 10% (v/v) heat inactivated FCS. IL-5 was detected colorimetrically using biotin-labeled antibodies followed by peroxidase-conjugated strepto-avidin and then peroxidase-specific colorimetric substrate. Standard curves were generated using known amounts of recombinant IL-5.

5 The detection limit of these assays was 40 pg/ml. All assays were performed in duplicate.

#### 9) IgM Secretion Assay

PBMC ( $3 \times 10^5$ /well) were cultured as described above with ODNs (1.5 µg/ml) for 72 hr. After this, supernatants were collected and IgM assayed by ELISA.

10 Briefly, 96 microtiter plates (NUNC, Denmark) were coated with anti-IgM antibodies and blocked with RPMI 1640 media. IgM was detected colorimetrically using peroxidase-labeled antibodies followed by peroxidase-specific colorimetric substrate. Standard curves were generated using known amounts of purified IgM. The detection limit of these assays was 50 ng/ml. All  
15 assays were performed in duplicate.

#### 10) Flow cytometry

Staining of surface antigens was performed as described (J. Fló and E. Massouh. Age-related changes of native and memory CD4 rat lymphocyte subsets in mucosal and systemic lymphoid organs. Developmental and  
20 comparative Immunology 21: 443-453, 1997). Anti CD19 (Clone LT19), CD86 (Clone BU63), CD40 (clone LOB 7/6), CD4 (clone S 3.5), MHC class I (Clone W6/32), MHC class II (Clone WR 18), anti CD4 (Clone S3.5) and anti CD11c

(Clone BU15) antibodies were purchased from Serotec (Raleigh, NC, USA). Apoptosis was estimated using the ANNEXIN V:FITC assay kit from Serotec (Raleigh, NC, USA).

Flow cytometric data of 10,000 cells/sample were acquired on a FACScan  
5 (Becton Dickinson Immunocytometry Systems, San Jose, CA). Data were analyzed using the computer program Win MDI, 2.8 , Interface Flow Cytometry Application (Joseph Trotter Copyright 1993-1998).

11) Immunization of monkeys against hepatitis B surface antigen (HBsAg)  
10 and evaluation of the humoral response

Twelve monkeys of the species *Cebus Apella* (2.5-3.5 Kg) were immunized with a pediatric dose of AgB (Pablo Cassará, Buenos Aires, Argentina) containing 10 µg of HBsAg adsorbed to alumina (25 mg of Al <sup>3+</sup>/ mg of HBsAg). This was administered alone (n= 3, 1 female, 2 males) or combined  
15 with indicated phosphorothioate ODN (150 µg/dose) (n=3, 1 female, 2 males). All vaccines were administered intramuscularly (i.m.) in the quadriceps muscle in a total volume of 1 ml. Monkeys were kept in the animal facility of the CEMIC (Centro Médico de Investigaciones Clínicas), Buenos Aires, Argentina. Animals were monitored daily by specialists and weighed once a  
20 week. Plasma was recovered by intravenous (i.v.) puncture before and at various times after immunization and immediately assayed for antibodies

using the commercial kit AUSAB (Abbot Laboratories, Illinois, USA). Titers are expressed in milliinternational units per ml.

## EXAMPLE 2

### 5 Selection of oligonucleotide sequences

WO 96/02555 and US 6,239,116 patents teach that to be immunostimulatory, oligonucleotides require sequences containing unmethylated CpG motifs (WO 96/02555, col. 13 lines 19-20 and US 6,239,116, col. 6 lines 1-3). EP 0 468 520 teaches that to be  
10 immunostimulatory, oligonucleotides require a palindromic sequence of at least 6 nucleotides long to be satisfactory (EP 0 468 520, col. 11, lines 34-37). Therefore, several non-CpG oligonucleotides, without palindromic sequences of at least 6 nucleotides long were proved using proliferation assays, cell differentiation assays, cytokine IL6 secretion assays and IgM  
15 secretion assays performed on human peripheral blood mononuclear leukocytes (PBMC). As a positive control, the CpG oligonucleotide 2006 of composition :

5' TCGTCGTTTTGTCGTTTTGTCGTT 3'

and phosphorothioate bonds described by Hartman and Krieg (Hartmann, G.,  
20 Krieg, A.M. 2000. Mechanism and function of a newly identified CpG DNA motif in human primary B cells. J. Immunol. 164:944.) was used. As a background control, either the phosphorothioate oligonucleotide IMT 023

(SEQ ID N<sup>o</sup>9) or IMT 022 (SEQ ID N<sup>o</sup>8) with very low activity on human cells was used.

Also as a "presumably" negative control an oligonucleotide with the same composition of the oligonucleotide 2006 but in which all the CpG dinucleotides have been replaced by GpC dinucleotides:

5' TGCTGCTTTTGTGCTTTTGTGCTT 3'

was used. This oligonucleotide was named ODN IMT 021 (SEQ ID N<sup>o</sup> 1).

Fig. 1 shows a proliferation assay performed with the above described oligonucleotides.

It was found, surprisingly, that the non CpG oligonucleotide IMT 021 was as active as the 2006 CpG oligonucleotide in proliferation assays (Fig. 1 and Table 1) if used at 1.5 µg/ml and approximately 40-60% active if used at 0.375 µg /ml.

**TABLE 1**

Induction of peripheral white blood cell proliferation by Phosphorothioate CpG (2006) and non-CpG oligonucleotides (IMT 021)

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX					
		(ODN at 1.5 µg/ml)			(ODN at 0.375 µg/ml)		
		Avg.	N	SD	Avg.	N	SD
2006	TCGTCGTTTTGTCGTTTTGTCGTT	17.95	4	2.39	13.83	14	1.55
IMT 021	TGCTGCTTTTGTGCTTTTGTGCTT	17.41	5	2.74	8.02	15	1.21

---

IMT 021 (SEQ. ID N° 1)

Avg.: Average; N: Number of Data; SD: Standard Deviation; ODN(S): phosphorothioate ODN

5 Immunostimulation was also evaluated by Flow Cytometry using the CD 19 general marker for human B cells and the CD 86 activation marker for human B cells.

Fig. 2 shows the activation of human B cells incubated with the non CpG ODN (S) IMT 021 as compared to the activation induced by incubation with  
10 the CpG ODN (S) 2006. As can be observed, both ODNs show similar activation patterns as compared with the ODN(S) IMT 023 background control when used at 1.5 ug/ml.

IL6 secretion was also evaluated in supernatants of PBMC incubated with the CpG ODN (S) 2006 and the non CpG ODN (S) IMT 021 (Table 2).

15

**TABLE 2**

Induction of human IL-6 secretion by phosphorothioate

CpG and non-CpG oligonucleotides 2006 and IMT 021

ODN (S)	<u>IL-6 (pg/ml)</u>			
	<u>expt. 1</u>	<u>expt. 2</u>	<u>expt. 3</u>	<u>expt. 4</u>
20 cells alone	0	0	0	0
2006	568	159	597	374

IMT 021 348 384 836 596

---

IMT 021 (SEQ. ID N<sup>o</sup> 1) . ODNs at 1.5 ug/ml

- 5 Results of this assay also indicate that the non-CpG ODN (S) IMT 021 is an effective immunostimulant when used at 1.5 ug/ml. Therefore, other non-CpG sequence variants of the 2006 oligonucleotide were investigated. Table 3 shows the results for six of these variants in which, the Cs or Gs of all the CpGs of this ODN were replaced by other nucleotides. As can be observed,
- 10 the Gs of the CpGs in the ODN(S) 2006 are not necessary for B cell proliferation or IL6 secretion. However, modification of the Cs of the CpGs is detrimental if the replacement is for As or Gs but not if it is for Ts. These results clearly indicate that stimulation of the B cell proliferation and IL6 secretion by the ODN(S) 2006 *is not at all associated with integrity of the*
- 15 *CpG group.*

**TABLE 3**

Induction of peripheral white blood cell proliferation and IL6 secretion by non-CpG variants of the ODN(S) 2006 oligonucleotide

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX			IL-6 (pg/ml)		
		(ODN at 0.375 $\mu$ g/ml)			(ODN at 6.0 $\mu$ g/ml)		
		Avg.	N	SD	Avg.	N	SD
20 2006 83.4	TCGTCGTTTTGTCGTTTTGTCGTT	13.83	14	1.55	215.8	4	



IMT 504 39.5	TCATCATTTCATTCATTTCATTCATT	13.08	2	3.80	285.2	2
IMT 505 3.1	TCCTCCTTTTGTCCCTTTTGTCCCTT	12.98	2	4.73	218.9	2
IMT 506 10	TCTTCCTTTTGTCTTTTGTCTTT 161.0	11.66	4	2.77	228.9	4
IMT 501 54.0	TAGTAGTTTGTAGTTTGTAGTT	5.42	2	2.03	66.8	2
IMT 502 91.4	TGGTGGTTTGTGGTTTGTGGTT	7.16	2	3.24	89.7	2
IMT 503	TTGTTGTTTTGTTGTTTTGTTGTT 215.5	9.87	2	1.90	334.9	2
<hr/>						
20 IMT 504 (SEQ ID N°2); IMT 505 (SEQ ID N°3); IMT 506 (SEQ ID N°4); IMT 501 (SEQ ID N°5); IMT 502 (SEQ ID N°6); IMT 503 (SEQ ID N°7)						

## EXAMPLE 3

25      Effect of structure modification on the immunostimulatory activity of  
non-CpG oligonucleotides: definition of the active motif.

In order to study the influence of the primary structure on the immunostimulatory activity of the non-CpG ODNs, several variants of the 2006 and IMT 021

30      oligonucleotides were synthesized. Table 4 shows the primary structure of some of the IMT 021 variants and the results of a proliferation and IL-6 assays performed in order to evaluate its immunostimulatory activity.

The ODN(S) IMT 021 contains T in positions 7,8,9,10,12,15,16,17,18,20,23 and 24. Replacement of these Ts with As (ODN IMT 022) or Cs (ODN IMT

023) results in a very significant loss of activity (about 76% and 75% respectively) in the proliferation assay and also in the IL-6 secretion assay (84% and 88% respectively). These results indicate that some or all the Ts in positions 7,8,9,10,12,15,16,17,18,20,23 and 24 are critical for the ODN IMT  
 5 021 immunostimulatory activity.

TABLE 4

Induction of peripheral white blood cell proliferation and IL-6 secretion by  
Phosphorothioate non-CpG oligonucleotides derived from the  
immunostimulatory IMT 021 oligonucleotide

10

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX (ODN at 0.375 $\mu$ g/ml)			IL-6 (pg/ml) (ODN at 6.0 $\mu$ g/ml)		
		Avg.	N	SD	Avg.	N	SD
IMT 021	TGCTGCTTTTGTGCTTTTGTGCTT	8.02	15	1.21	181.3	4	49.9
IMT 022	TGCTGCAAAAGAGCAAAAGAGCAA	1.24	2	0.30	0.0	2	
0.0							
IMT 023	TGCTGCCCCCGCGCCCCGCGCCC	1.07	4	0.12	0.0	3	
0.0							
IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 023 (SEQ. ID N° 9)							

20

In order to investigate the composition of the active center (motif) of the non-CpG oligonucleotides, several ODNs were assayed at 0.375  $\mu$ g/ml (Fig. 3). It was found that:

- 25 a) The presence of TG groups in an oligonucleotide is not at all sufficient  
 (in disagreement with WO 01-22972) for a strong immunostimulation.

For example, ODNs IMT 028, IMT 046, IMT 546, IMT 547 and IMT 550, all of which have two TGs, have very poor activity. On the other hand, ODN IMT 543 that does not possess a TG dinucleotide has the same (low) activity as ODN IMT544, which is an ODN with the same general composition, but which contains two TG dinucleotides.

b) The presence of the tetranucleotide TTTT in an oligonucleotide is not at all sufficient (in disagreement with WO 01-22972) for a strong immunostimulation. For example, ODNs IMT 034, IMT 057, IMT 028, IMT 543 and IMT 544, which have two TTTT motifs, have very poor activity.

c) The presence of more than 25% of Ts is not at all sufficient (in disagreement with WO 01-22972) for a strong immunostimulation. For example, ODNs IMT 034(50% T), IMT 057 (71% T), IMT 028 (33% T), IMT 543(58% T), IMT 544 (50% T), IMT 059 (83% T), IMT 040 (42% T), IMT 046 (33% T), IMT 545 (42% T), IMT 550 (33% T) and IMT 546(42% T), all of which have more than 25% of T, have very poor activity.

d) The presence of more than 25% of Ts plus the presence of TG dinucleotides, is not at all sufficient (in disagreement with WO 01-22972) for strong stimulation. For example, ODN IMT 028, IMT 046, IMT 544 and IMT 546, which have more than 25% of Ts and two TGs, have very poor activity.

- 5 e) The presence of more than 25% of Ts plus the presence of the tetranucleotide TTTT, is not at all sufficient (in disagreement with WO 01-22972) for strong stimulation. For example, ODN IMT 059, IMT 034, IMT 057, IMT 028, IMT 543 and IMT 544, which have more than 25% of Ts and two TTTT, have very poor activity.
- f) The presence of TG dinucleotides plus the presence of the tetranucleotide TTTT is not at all sufficient (in disagreement with WO 01-22972) for strong stimulation. For example, ODN IMT 028 and IMT 544, which have two TGs and two TTTT, have very poor activity.
- 10 g) The presence of TG dinucleotides plus the presence of the tetranucleotide TTTT, plus the presence of more than 25% Ts, is not at all sufficient (in disagreement with WO 01-22972) for strong stimulation. For example, ODN IMT 028 and IMT 546, which have two TGs, two TTTT and more than 25% Ts, have very poor activity.
- 15 h) Some Ts may be relevant for the activity of non CpG ODNs, but they are not necessarily related to a TG dinucleotide or to a TTTT motif. For example, the replacement of 2 Ts in ODN IMT 540, which are not involved in any TG or TTTT motif, is strongly detrimental (ODN 545). It is worth noting that both, ODN IMT 540 and IMT 545 have more than
- 20 25% T and do not have any TTTT or TG motifs. It is also worth noting that the substitution, in the poorly immunostimulatory ODN IMT 545 of two Ts, that generates two TGs dinucleotides, two TTTT motifs and

increases the total content of Ts (ODN IMT 544) does not increase the activity (in disagreement with WO 01-22972).

- 5 i) The presence of the tetranucleotide CCCC and/or more than 50% of Cs is not at all sufficient (in disagreement with WO 01-22972) for a strong immunostimulation. For example, ODNs IMT 023, which have two CCCC and 63% of C, have very poor activity.
- j) Replacement of some Ts for Cs or Gs in some T-rich oligonucleotides increases the activity (in disagreement with WO 01-22972). For example, replacement of 3 Ts in ODN IMT 059 by Cs (ODN IMT 037)  
10 increases the activity in a statistically significant manner. Therefore, the presence of Cs combined with some Ts may be relevant to achieve strong immunostimulation.
- k) In general, T-rich oligonucleotides may have some immunostimulatory activity in a phosphorothioate backbone, but are inactive in a  
15 phosphodiester backbone (in disagreement with WO 01-22972). For example, Table 5 shows that IMT 053, which consists of a polyT chain 24 nucleotides long, is able to stimulate the secretion of IL6 in a phosphothioate backbone, but not in a phosphodiester backbone. The same is true for ODN IMT 021, which contains more than 25% T and 6  
20 TG dinucleotides. These two ODNs are also unable, in a phosphodiester backbone, to stimulate the expression of the

activation markers CD 86, CD 40 and MHC class I on CD 19 positive (B) cells (Fig. 4).

- 5 I) In general, immunostimulatory oligonucleotides in a phosphodiester backbone are inactive as a double strand (in disagreement with WO 01-22972). For example, the strong stimulatory ODNs 2006 and IMT 504 are inactive as a double strand (not shown). Similar results were described in mice in the following publication: Zelenay, S., Elias, F. and Flo, J. Eur. J. Immunol. 33: 1382-92 (2003).

- 10 Taking into account all these facts, it was investigated which one is the minimal sequence necessary and sufficient that should be present in a non-CpG oligonucleotide in order to obtain a strong immunostimulatory activity, in a phosphodiester (natural) as well as in modified backbones.

15

**Table 5**

Induction of IL6 secretion by phosphorothioate or phosphodiester

PyNTTTTGT ODNs

ODN (O)	Sequence	IL-6 Secretion (pg/ml)		
		Avg.	N	SD
		38	4	17
IMT 022 (S)	TGCTGCAAAAGAGCAAAAGAGCAA	27	4	12
IMT 022 (O)	TGCTGCAAAAGAGCAAAAGAGCAA	123	4	21
IMT 053 (S)	TTTTTTTTTTTTTTTTTTTTTTTTTT	24	4	24
IMT 053 (O)	TTTTTTTTTTTTTTTTTTTTTTTTTT			

IMT 021 (S)	TGCTGCTTTTGTGCTTTTGTGCTT	182	4	51
IMT 021 (O)	TGCTGCTTTTGTGCTTTTGTGCTT	35	4	12
IMT 504 (S)	TCATCATTTTGTCAATTTTGTCAAT	285	4	39
IMT 504 (O)	TCATCATTTTGTCAATTTTGTCAAT	210	4	43
IMT 506 (S)	TCTTCTTTTGTCTTTTGTCTTT	228	4	70
IMT 506 (O)	TCTTCTTTTGTCTTTTGTCTTT	250	4	32
2006 (S)	TCGTCGTTTTGTCGTTTTGTCGTT	216	4	84
2006 (O)	TCGTCGTTTTGTCGTTTTGTCGTT	228	4	40
None	---	28	4	14

IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 053 (SEQ. ID N° 19); IMT 504 (SEQ. ID N° 2); IMT 506 (SEQ. ID N° 4)

(S): phosphorothioate. (O): phosphodiester. Phosphorothioate ODNs were added (6 µg/ml) at time 0 of incubation.

- 5 Phosphodiester ODNs were added as follows: 30 µg/ml at time 0, 30 µg/ml after 4 hr and 30 µg/ml after 16 hr of incubation.

The analysis of hundreds of ODNs (not shown) allowed definition of a core  
 10 sequence (*motif*) responsible for the strong immunostimulatory activity of non-  
 CpG ODNs as measured by B cell proliferation and IL6 secretion. This *motif*  
 is the following:



- wherein  $X_1$  is C or T  
 15 wherein  $X_2$  is C, T, G or A;  
 wherein  $X_3$  is T or A;  
 wherein  $X_4$  is T, C or G;  
 wherein  $X_5$  is T, C or G;  
 wherein  $X_6$  is T or G;  
 20 wherein  $X_7$  is G;  
 wherein  $X_8$  is T  
 wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are Ts

Table 6 shows the effect of changes in each of the nucleotides of the two non-CpG motifs present in the ODN IMT 504 (motif: CATTTTGT). Replacement of the C in position 1 of the motif by A (ODN IMT 531) or G (ODN IMT 533) resulted in a loss of about 50% in the activity. However, replacement of this C by T (ODN IMT 532) did not change the activity. Thus, in order to obtain maximal activity the first position of the motif should be occupied by a pyrimidine nucleotide (C or T). In position 2 of the motif, A (ODN IMT 504) or T (ODN IMT 535) or C (ODN IMT 534) are equivalent options.

10 In order to study the influence of a G nucleotide in position 2 of the motif without introduction of a CpG dinucleotide, ODNs with a T in the first position of the motif were synthesized (Table 7). As can be observed, any nucleotide in the second position of the motif is equivalent.

Replacement of the G in position 7 of the motif by A (ODN IMT 541), T (ODN IMT 542) or C (ODN IMT 543) is detrimental (Table 5). Thus, in order to obtain maximal activity position 7 of the motif should be preferably G.

Regarding the positions of the motif occupied by Ts, the most sensitive is the one in position 8. Replacement of this T for A (ODN IMT543) or C (ODN IMT599) or G (ODN IMT 604) is detrimental. Thus, in order to obtain maximal activity, position 8 of the motif should be preferably T. In position 3 of the motif the T could be replaced by A (ODN IMT IMT537) without activity loss. However, replacement of this T by C (ODN IMT595) or G (ODN IMT600) is



detrimental. Thus, in order to obtain maximal activity, position 3 of the motif should be preferably T or A. In position 4 of the motif the T could be replaced by C (ODN IMT IMT596) or G (ODN IMT601) without significant activity loss. However, replacement of this T by A (ODN IMT538) is detrimental. Thus, in order to obtain maximal activity, position 4 of the motif should be preferably T or C or G. In position 5 of the motif, the T could be replaced by C (ODN IMT IMT597) or G (ODN IMT602) without significant activity loss. However, replacement of this T by A (ODN IMT539) is detrimental. Thus, in order to obtain maximal activity, position 5 of the motif should be preferably T or C or G. In position 6 of the motif, the T could be replaced by G (ODN IMT IMT603) without significant activity loss. However, replacement of this T by A (ODN IMT540) is detrimental. Thus, in order to obtain maximal activity, position 6 of the motif should be preferably T or G.

Simultaneous replacement of T by A in position 3 of the motif and T by G in position 6 of the motif did not result in any loss of activity (ODN IMT 608). However, in this case and in order to obtain maximal activity, positions 4 and 5 should be occupied by Ts (see ODNs IMT609, IMT610, IMT611, IMT612 and IMT614).

Replacement of two or more of the Ts by As within the motif results in more than 70% loss of activity (ODNs IMT 545, IMT 546, IMT 547, IMT 548, IMT 549, IMT 550, IMT 551 and IMT 552).

Taking these results into consideration, it can also be concluded that in order to obtain a strong immunostimulatory activity, two or more of the positions 3,4,5 and 6 of the motif should be occupied by Ts.

TABLE 6

Induction of peripheral white blood cell proliferation by phosphorothioate non-CpG oligonucleotides of this invention derived from ODN IMT 504

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX (ODN at 0.375 µg/ml)		
		Avg.	N	SD
2006	TCGTCGTTTTGTCGTTTTGTCGTT	13.35	16	2.23
IMT 022	TGCTGCAAAAGAGCAAAAGAGCAA	1.34	24	0.52
IMT 504	TCATCATTGTCATTGTCATT	13.85	16	4.36
IMT 531	TCATAATTTGTAATTTGTCATT	7.59	12	2.71
IMT 532	TCATTATTTGTTATTTGTCATT	13.80	12	3.41
IMT 533	TCATGATTTGTGATTTGTCATT	6.20	12	2.71
IMT 534	TCATCCTTTGTCCTTTGTCATT	11.67	12	4.49
IMT 535	TCATCTTTTGTCTTTTGTGTCATT	12.49	12	2.51
IMT 537	TCATCAATTTGTCAATTTGTCATT	14.31	12	4.03
IMT 595	TCATCACTTTGTCACTTTGTCATT	9.16	15	2.30
IMT 600	TCATCAGTTTGTGAGTTTGTGTCATT	7.38	12	2.69
IMT 538	TCATCATATTGTCATATTGTCATT	7.80	16	3.38
IMT 596	TCATCATCTTGTGTCATTGTCATT	12.61	20	3.37
IMT 601	TCATCATGTTGTGTCATTGTCATT	11.47	8	1.35
IMT 539	TCATCATTATGTCATTATGTCATT	5.63	16	3.52
IMT 597	TCATCATTCTGTCATTCTGTCATT	13.98	20	3.78
IMT 602	TCATCATTGTGTCATTGTGTCATT	13.23	20	3.47
IMT 540	TCATCATTTAGTCATTAGTCATT	9.62	16	3.47
IMT 603	TCATCATTTGGTCATTGGTCATT	13.91	20	3.35
IMT 541	TCATCATTTTATCATTTTATCATT	9.30	12	2.20
IMT 542	TCATCATTTTTCATTTTTCATT	9.78	12	2.60
IMT 543	TCATCATTTTCTCATTTTCTCATT	4.75	12	2.25
IMT 544	TCATCATTTTGACATTTGACATT	4.27	12	1.30
IMT 599	TCATCATTTTGCCATTTGCCATT	5.39	16	1.98
IMT 604	TCATCATTTTGGCATTTGGCATT	7.60	16	2.63
IMT 608	TCATCAATTGGTCAATTGGTCATT	12.98	12	2.22
IMT 609	TCATCAAATGGTCAAATGGTCATT	8.58	12	1.53

IMT 610	TCATCAACTGGTCAACTGGTCATT	9.07	12	1.95
IMT 611	TCATCAAGTGGTCAAGTGGTCATT	5.97	12	1.79
IMT 612	TCATCAATAGGTCAATAGGTTCATT	3.15	12	0.75
IMT 614	TCATCAATGGGTCAATGGGTTCATT	5.18	12	1.54
IMT 545	TCATCATTAGACATTAGACATT	4.19	12	1.81
IMT 546	TCATCATTATGACATTATGACATT	2.90	12	1.16
IMT 547	TCATCATATTGACATATTGACATT	3.34	12	1.16
IMT 548	TCATCATTAAGACATTAAGACATT	3.40	12	1.53
IMT 549	TCATCATATAGACATATAGACATT	3.51	12	1.03
IMT 550	TCATCATAATGACATAATGACATT	1.91	12	0.71
IMT 551	TCATCATAAAGACATAAAGACATT	2.17	12	0.72
IMT 552	TCATCAAAAAGACAAAAAGACATT	1.37	12	0.83

IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2); IMT 531 (SEQ. ID N° 14); IMT 532 (SEQ. ID N° 15); IMT 533 (SEQ. ID N° 16); IMT 533 (SEQ. ID N° 17); IMT 534 (SEQ. ID N° 18); IMT 535 (SEQ. ID N° 19); IMT 537 (SEQ. ID N° 30); IMT 538 (SEQ. ID N° 31); IMT 539 (SEQ. ID N° 32); IMT 540 (SEQ. ID N° 33); IMT 541 (SEQ. ID N° 34); IMT 542 (SEQ. ID N° 35); IMT 543 (SEQ. ID N° 36); IMT 544 (SEQ. ID N° 37); IMT 545 (SEQ. ID N° 38); IMT 546 (SEQ. ID N° 39); IMT 547 (SEQ. ID N° 40); IMT 548 (SEQ. ID N° 41); IMT 549 (SEQ. ID N° 42); IMT 550 (SEQ. ID N° 43); IMT 551 (SEQ. ID N° 44); IMT 552 (SEQ. ID N° 45); IMT 539 (SEQ. ID N° 82); IMT 597 (SEQ. ID N° 83); IMT 602 (SEQ. ID N° 84); IMT 540 (SEQ. ID N° 85); IMT 603 (SEQ. ID N° 87); IMT 544 (SEQ. ID N° 88); IMT 599 (SEQ. ID N° 89); IMT 604 (SEQ. ID N° 90); IMT 608 (SEQ. ID N° 75); IMT 609 (SEQ. ID N° 76); IMT 610 (SEQ. ID N° 77); IMT 611 (SEQ. ID N° 78); IMT 612 (SEQ. ID N° 79); IMT 614 (SEQ. ID N° 81).

TABLE 7

Induction of peripheric white blood cell proliferation and IL-6 secretion by

Phosphorothioate non-CpG oligonucleotides with changes in the second

position of the immunostimulatory motif

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX			IL-6 (pg/ml)		
		(ODN at 0.375 µg/ml)			(ODN at 6.0 µg/ml)		
		Avg.	N	SD	Avg.	N	SD
IMT 532	TCATTATTTTGTATTTTGTTCATT	10.10	4	1.61	221,7	4	35,3
IMT 197	TCATTTTTTTGTTTTTTGTTCATT	10.84	4	1.06	386,1	4	123,3
IMT 198	TCATTGTTTTGTGTTTTTGTTCATT	10.00	4	1.75	235,3	4	27,1
IMT 199	TCATTCTTTTGTCTTTTGTTCATT	12.36	3	1.71	235,0	4	6,0

IMT 532 (SEQ. ID N° 15); IMT 197 (SEQ. ID N° 10); IMT 198 (SEQ. ID N° 11); IMT 199 (SEQ. ID N° 12)

To investigate the effect of changes in the position of the motif within the ODN chain, the non-CpG motif CATT TGT present in ODN IMT504 was introduced into different locations in a 24-nucleotide long poly T chain (Fig. 5). As can be seen, the poly T ODN (S) by itself had a significant mitogenic activity. On the other hand, introduction of only one CATT TGT motif resulted in an increment of 1.2 to 1.8 times (measured by proliferation assays), or 2.4 to 3.5 times (measured by IL-6 secretion assays) in the immunostimulatory activity of the polyT chain, depending on where the motif was located. A distance of at least two nucleotides from the 5' end or four from the 3' end seems to be necessary in order to reach maximal activity (ODNs IMT174 to IMT 179). Introduction of two motifs in optimal positions (ODN IMT182) resulted in an increment of 1.9 times (measured by proliferation assays), or 2.9 times (measured by IL-6 secretion assays) in the immunostimulatory activity of the polyT chain. This indicates that the contribution of the second motif is negligible. On the other hand, the activity of the most effective ODNs like ODN IMT504 is more than 3 times larger than the activity of the poly T as measured by proliferation assays, a fact which suggests that there is also a significant influence of the composition of at least some of the nucleotides surrounding the motif of this invention on the overall activity of the ODN.

20

#### EXAMPLE 4

##### Effect of structure modification on the immunostimulatory activity of

non-CpG oligonucleotides of this invention: influence of the nucleotide  
composition outside of the active motif.

Results shown in Fig. 5 indicate that composition of the oligonucleotide outside of the non-CpG core motif is important in order to reach optimal immunostimulatory activity. Therefore, a number of oligonucleotides were synthesized with changes in the composition of the nucleotides surrounding the motifs. Table 8 shows the effect of changes in the composition of the first four nucleotides of the 5' end and of the last four nucleotides of the 3' end in the immunostimulatory activity of ODN IMT 504. As can be observed, the best choices are: C or G in position -1 respect to the two motifs, C, T or G in position -2, any nucleotide in positions -3 and -4, A or T in position +1, G in position +2, any nucleotide in positions +3 and G in position +4.

Of course, other nucleotide combinations not represented in this table may have equal or even better effect on the activity of the non-CpG immunostimulatory oligonucleotides. One of ordinary skill in the art can empirically determine other effective combinations.

**TABLE 8**

Induction of peripheral white blood cell proliferation by non- CpG ODN(S) IMT  
504 with variations in the composition outside of the two immunostimulatory  
motifs

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX			IL-6 (pg/ml)		
		(ODN at 0.375 µg/ml)			(ODN at 6.0 µg/ml)		
		Avg.	N	SD	Avg.	N	SD
IMT 552	TGCTGCAAAAGAGCAAAAGAGCAA	1,9	12	0,7	<39,9	4	—
IMT 504	TCATCATTTTGTCAATTTGTCAAT	11,0	8	0,9	171	2	14
IMT 559	ACATCATTTTGTCAATTTGTCAAT	13,4	4	1,0	128	3	3
IMT 560	CCATCATTTTGTCAATTTGTCAAT	12,1	4	0,9	145	4	32
IMT 561	GCATCATTTTGTCAATTTGTCAAT	9,5	4	1,1	209	4	74
IMT 562	TAATCATTTTGTCAATTTGTCAAT	12,4	4	1,5	171	4	24
IMT 563	TTATCATTTTGTCAATTTGTCAAT	11,1	3	0,7	172	4	63
IMT 564	TGATCATTTTGTCAATTTGTCAAT	12,8	4	0,9	118	4	38
IMT 565	TCCTCATTTTGTCAATTTGTCAAT	14,1	4	0,5	135	4	25
IMT 566	TCCTCATTTTGTCAATTTGTCAAT	13,9	4	2,2	157	4	28
IMT 567	TCGTCAATTTTGTCAATTTGTCAAT	12,9	4	0,7	259	4	25
IMT 568	TCAACATTTTGTCAATTTGTCAAT	10,3	4	1,1	153	4	35
IMT 569	TCACCATTTTGTCAATTTGTCAAT	15,0	4	1,2	199	3	12
IMT 570	TCAGCATTTTGTCAATTTGTCAAT	12,5	4	0,4	181	3	2
IMT 571	TCATCATTTTGTCAATTTGTAAAT	14,2	4	0,7	163	4	26
IMT 572	TCATCATTTTGTCAATTTGTATAT	13,2	4	1,4	182	3	67
IMT 573	TCATCATTTTGTCAATTTGTGATT	8,7	4	0,7	177	4	29
IMT 574	TCATCATTTTGTCAATTTGTCTTT	8,1	4	0,7	179	3	58
IMT 575	TCATCATTTTGTCAATTTGTCTTT	10,8	4	0,8	150	3	47
IMT 576	TCATCATTTTGTCAATTTGTCTTT	12,6	4	1,1	202	4	55
IMT 577	TCATCATTTTGTCAATTTGTCAAT	10,0	4	1,4	243	3	8
IMT 578	TCATCATTTTGTCAATTTGTCACT	10,9	4	0,9	213	2	9
IMT 579	TCATCATTTTGTCAATTTGTCACT	10,7	4	1,7	182	4	18
IMT 580	TCATCATTTTGTCAATTTGTCACT	11,7	4	1,4	194	4	36
IMT 581	TCATCATTTTGTCAATTTGTCACT	10,7	4	1,0	173	4	18
IMT 582	TCATCATTTTGTCAATTTGTCACT	12,1	4	0,8	242	3	231

IMT 552 (SEQ. ID N° 45); IMT 504 (SEQ. ID N° 2); IMT 559 (SEQ. ID N° 46); IMT 560 (SEQ. ID N° 47); IMT 561 (SEQ. ID N° 48); IMT 562 (SEQ. ID N° 49); IMT 563 (SEQ. ID N° 50); IMT 564 (SEQ. ID N° 51); IMT 565 (SEQ. ID N° 52); IMT 566 (SEQ. ID N° 53); IMT 567 (SEQ. ID N° 54); IMT 568 (SEQ. ID N° 55); IMT 569 (SEQ. ID N° 56); IMT 570 (SEQ. ID N° 57); IMT 571 (SEQ. ID N° 58); IMT 572 (SEQ. ID N° 59); IMT 573 (SEQ. ID N° 60); IMT 574 (SEQ. ID N° 61); IMT 575 (SEQ. ID N° 62); IMT 576 (SEQ. ID N° 63); IMT 577 (SEQ. ID N° 64); IMT 578 (SEQ. ID N° 65); IMT 578 (SEQ. ID N° 66); IMT 580 (SEQ. ID N° 67); IMT 581 (SEQ. ID N° 68); IMT 582 (SEQ. ID N° 69)

## EXAMPLE 5

### Effect of structure modification on the immunostimulatory activity of

10 non-CpG oligonucleotides of this invention: influence of the size of the  
oligonucleotide

Table 9 shows that ODNs with one immunostimulatory motif are active if the chain is 16 or more nucleotides long. Activity is maximal if the ODN is between 20 and 25 nucleotides long. *All the ODNs of within this range bearing at least one non-CpG motif of this invention assayed during this study (more than hundred) were very active.*

TABLE 9

Effect of the size of phosphorothioate non-CpG oligonucleotides on B cells proliferation and IL6 secretion

10

ODN (S)	SEQUENCE (5'-3')	PROLIFERATION INDEX (ODN at 0.375 $\mu$ g/ml)			IL-6 (pg/ml) (ODN at 6.0 $\mu$ g/ml)		
		Avg.	N	SD	Avg.	N	SD
IMT 187	TTTTCATTTTGT	0.56	8	0.20	<50	4	—
IMT 188	TTTTCATTTTGTTTT	2.76	8	1.73	147	4	40
IMT 189	TTTTCATTTTGTTTTTTTT	8.21	8	3.19	227	4	21
IMT 175	TTTTCATTTTGTTTTTTTTTTTT	8.92	8	1.80	220	4	18
IMT 179	TTTTTTTTTTTCATTTTGTTTT	6.06	8	2.00	224	4	82
IMT 191	TTTTCATTTTGTTTTTTTTTTTTTTTT	6.64	8	2.37	205	4	15

IMT 187 (SEQ ID N°70); IMT 188 (SEQ ID N°71); IMT 189 (SEQ ID N°72); IMT 179 (SEQ ID N°73); IMT 191 (SEQ ID N°77)

## EXAMPLE 6

15 Induction of peripheral white blood cell IgM secretion by phosphorothioate non-CpG oligonucleotides of this invention

Induction of IgM secretion in peripheral white blood cells is another important marker of immunostimulatory activity of oligonucleotides. Table 10 shows the

stimulation of IgM secretion by several of the phosphorothioate non-CpG oligonucleotides described above. As can be observed, the most active non-CpG ODN(S)s in induction of proliferation and IL6 secretion are also the best in induction of IgM secretion.

TABLE 10

Induction of peripheral white blood cell IgM secretion by phosphorothioate non-CpG oligonucleotides of this invention

ODN (S)	SEQUENCE (5'-3')	IgM SECRETION (ng/ml) (ODN at 1.50 µg/ml)		
		Avg.	N	SD
cells alone		76	9	50
2006	TCGTCGTTTTGTCGTTTTGTCGTT	778	47	204
IMT 503	TTGTTGTTTTGTTGTTTTGTTGTT	640	19	220
IMT 504	TCATCATTTTGTCAATTTGTCAAT	912	24	244
IMT 505	TCCTCCTTTTGTCTTTTGTCTTT	664	23	336
IMT 506	TCTTCTTTTGTCTTTTGTCTTT	751	43	237
IMT 509	AAAAAACTAAAAAACTAAAAAA	195	17	77

IMT 504 (SEQ ID N°2); IMT 505 (SEQ ID N°3); IMT 506 (SEQ ID N°4); IMT 503 (SEQ ID N°7); IMT 509 (SEQ ID N°13)

## EXAMPLE 7

Stimulation of the expression of CD40, MHC I and MHC II in B lymphocytes by phosphorothioate non-CpG oligonucleotides of this invention



Human PMBC were incubated with ODN IMT 504, ODN 2006 as a positive control and ODN IMT 022 as a negative control (Fig 6). As can be observed, ODN IMT 504 is as active as the CpG ODN 2006 for stimulation of the expression of CD40, MHC I and MHC II on B lymphocytes.

5

## EXAMPLE 8

Stimulation of purified B lymphocytes by non-CpG oligonucleotides of this invention

Human CD19<sup>+</sup> (B) cells were purified to more than 95% purity. Table 11 and Fig. 7 show that the immunostimulatory activity on these purified cells is comparable to the one observed using human PMBC. These results indicate that stimulation by the non-CpG oligonucleotides of this invention on human cells is direct.

10

TABLE 11

Induction of proliferation, IL6 and IgM secretion on purified B cells by phosphorothioate non-CpG oligonucleotides of this invention

ODN (S)	SEQUENCE (5'-3')	Proliferation Index (ODN at 0.375 $\mu$ g/ml)			IL-6 (pg/ml) (ODN at 6.0 $\mu$ g/ml)			IgM (ng/ml) (ODN at 1.5 $\mu$ g/ml)		
		Ave.	N	SD	Ave.	N	SD	Ave.	N	SD
IMT 022	TGCTGCAAAAGAGCAAAAGAGCAA	1,28	16	0,80	1019	5	665	469	7	182
IMT 504	TCATCATTTTGTTCATTTTGTTCATT	47,95	16	6,41	11002	5	884	1274	9	252
IMT 506	TCTTCCTTTTGTCTTTTGTCTTT	25,58	16	7,84	10933	3	1628	1083	10	430
2006	TCGTCGTTTTGTCTGTTTTGTCTGTT	60,82	24	9,79	7631	6	997	1292	12	245

IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N°2); IMT 506 (SEQ. ID N°4)

## EXAMPLE 9

Immunostimulation by phosphodiester non-CpG oligonucleotides of this invention

5 Fig 8 and Table 12 show the effect of phosphodiester non-CpG oligonucleotides of this invention on human PMBC. Since phosphodiester oligonucleotides are very sensitive to nucleases they were added to the culture three times (0, 4, and 16 hr) to a final concentration of 30 µg/ml. A very potent phosphodiester CpG ODN (ODN 2080) was used as positive control in cytometric assays (Hartmann G, Weeratna R, Ballas ZK, Payette P, Suparto, I, Rasmussen WL, Wadschmidt M, Sajuthi D, Purcells RH, Davis HL, Krieg AM. Delineation of a CpG phosphorothioate oligodeoxynucleotide for activating primate immune responses *in vitro* and *in vivo*. J. Immunol.164, 1617-1624, (2000)). As can be observed, under these conditions, 10 phosphodiester ODNs bearing non-CpG motifs of this invention have immunostimulatory activity. Regarding this, it is worth noting the differences of these non-CpG ODNs with those non-CpG ODNs claimed in WO 01/22972 (see table 5 and Fig 4).

## TABLE 12

20 Immune-stimulation by phosphodiester non-CpG oligonucleotides of this invention

ODN (O)	SEQUENCE (5'-3')	IL-6 Secretion (pg/ml)		
		(ODN at 6.0 µg/ml)		
		Avg.	N	SD
IMT 022	TGCTGCAAAAGAGCAAAAGAGCAA	23	8	10
IMT 053	TTTTTTTTTTTTTTTTTTTTTTTTTT	24	4	24
IMT 504	TCATCATTGTCATTGTCATT	207	6	46
IMT 506	TCTTCTTTTGTCTTTTGTCTTT	403	3	220
2006	TCGTCGTTTTGTCGTTTTGTCGTT	228	4	40

IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ ID N°2); IMT 053 (SEQ ID N°19); IMT 506 (SEQ ID N°4)

#### EXAMPLE 10

Induction of cell proliferation in peripheral white blood cells of monkeys of the  
species *Cebus apella* and *Macaca fascicularis* by non-CpG ODN(S)s of this  
invention

Non-CpG ODN(S)s of this invention are not very effective immunostimulants in mice, pigs and sheep. However, they are effective in monkeys. For example, Fig.9 shows the immunostimulatory activity of non-CpG ODN(S)s on peripheral white blood cells of monkeys of the species *Cebus apella* and *Macaca fascicularis*. According to these results the most effective non-CpG ODNs in non human primates are those bearing the CATTTTGT motif. Therefore, monkeys can be used as animal models for research on clinical applications of these non-CpG ODN(S)s.

#### EXAMPLE 11

### Vaccination of subjects

Vaccine formulations containing the non-CpG ODNs adjuvant of this invention can be used to vaccinate subjects against a variety of bacterial and viral disease agents and against tumor cells. Table 13 shows the effect of inoculation of monkeys of the species *Cebus apella* with a vaccine formulation that includes recombinant Hepatitis B surface antigen (rHBsAg) and alumina in presence or absence of ODN IMT 504 as adjuvant.

**TABLE 13**

Anti-HBs responses in *Cebus apella* immunized against HBsAg with IMT

#### ODNs

ODN (S)	Pre - prime	Post - prime		Post - boost
		14 days	28 days	14 days
None	0	0,4 ± 0,1	14 ± 20	190±52
IMT 504	0	10 ± 14	113 ± 114	659±296
2006	0	22 ± 24	67 ± 57	705±315

As can be observed, there is a dramatic increment in the title of anti-HBsAg in animals vaccinated with HBsAg plus IMT ODN(S) 504 as compared with those vaccinated with HBsAg alone. CpG ODN 2006 has a performance as adjuvant similar to the non-CpG ODNs of this invention.

In particular, a human can be vaccinated against hepatitis B by administration of a vaccine formulation that includes recombinant Hepatitis B surface

antigen (rHBsAg) and alumina and one or more of the oligonucleotides of this invention as adjuvants. The amount of rHBsAg in each dose, and the administration schedule, can vary according to the age of the human. For example, for humans from about birth to about 12 years old, a three-dose schedule of about 2.5  $\mu$ g to about 5  $\mu$ g of rHBsAg can be administered at 0, 1-3 months afterward and 4-18 months afterward, preferably at 0, 2 and 6 months. One or more of the oligonucleotides of this invention can be present in the formulation from about 10  $\mu$ g to about 1,000  $\mu$ g per dose.

For humans from about 12 to about 60 years old, a three-dose schedule of about 5  $\mu$ g to about 40  $\mu$ g of rHBsAg can be administered at 0, 1-3 months afterward and 4-18 months afterward, preferably at 0, 2 and 6 months. One or more of the oligonucleotides of this invention can be present in the formulation from about 10  $\mu$ g to about 1,000  $\mu$ g per dose.

#### EXAMPLE 12

##### Stimulation of the expression of costimulatory molecules in malignant B-lymphocytes by phosphorothioate non-CpG oligonucleotides

The stimulation of malignant B lymphocytes recovered from blood of patients suffering from chronic lymphocytic leukemia (CLL) by phosphorothioate non-CpG oligonucleotides of this invention was examined by flow cytometric (FACS) analysis. The results of a typical FACS analysis are shown in Fig. 10.

As can be seen, the phosphorothioate non-CpG oligonucleotide IMT 504 of this invention is able to stimulate expression of CD86, CD40 and MHC class I surface costimulatory molecules on malignant B lymphocytes so well as the CpG ODN 2006.

#### EXAMPLE 13

##### Stimulation of malignant B cell apoptosis by phosphorothioate non-CpG oligonucleotides

The stimulation of apoptosis in malignant B lymphocytes recovered from blood of patients suffering from chronic lymphocytic leukemia (CLL) by phosphorothioate non-CpG oligonucleotides of this invention was examined by flow cytometric (FACS) analysis using as apoptosis markers propidium iodide and annexin V. The result of a typical FACS analysis are shown in Fig 11. As can be observed, the non CpG oligonucleotide IMT 504 of this invention is able to increase the apoptosis of the leukemia cells so well as the CpG ODN 2006.

#### EXAMPLE 14

##### Stimulation of the expression of costimulatory molecules on plasmacytoid dendritic cells by phosphorothioate non-CpG oligonucleotides of this invention

The stimulation of plasmacytoid dendritic cells recovered from blood of normal donors by phosphorothioate non-CpG oligonucleotides of this invention was examined by flow cytometric (FACS) analysis. The results of a

typical FACS analysis are shown in Fig. 12. As can be seen, the phosphorothioate non-CpG oligonucleotide IMT 504 of this invention is able to stimulate expression of CD86, CD40 and MHC class I surface molecules on plasmacytoid dendritic cells so well as the CpG ODN 2006.

5

Table 14 shows that the non-CpG oligonucleotides of this invention are also able to stimulate purified human plasmacytoid dendritic cells to secrete interferon alpha if the oligonucleotide backbone is phosphodiester.

10

TABLE 14

Interferon alpha secretion by purified plasmacytoid dendritic cells incubated with phosphorothioate and phosphodiester non-CpG oligonucleotides of this invention

ODN	SEQUENCE (5'-3')	IFN Alpha Secretion (pg/ml)		
		Avg.	N	SD
IMT 022 (S)	TGCTGCAAAAGAGCAAAAGAGCAA	<31	3	—
IMT 504 (S)	TCATCATTTTGTTCATTTTGTTCATT	<31	3	—
IMT 022 (O)	TGCTGCAAAAGAGCAAAAGAGCAA	<31	3	—
IMT 504 (O)	TCATCATTTTGTTCATTTTGTTCATT	2300	3	632

15

IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ ID N°2). Phosphorothioate ODNs were added (1.5 µg/ml) at time 0 of incubation. Phosphodiester ODNs were added as follows: 30 µg/ml at time 0, 30 µg/ml after 4 hr and 30 µg/ml after 16 hr of incubation.

## EXAMPLE 15

Treatment of subjects with tumoral disease

Pharmaceutical formulations containing one or more of the immunostimulatory oligonucleotides of this invention, can be used to treat subjects against a variety of tumoral diseases. In particular, a human with a Melanoma can be treated by administration of a pharmaceutical formulation containing the oligonucleotide of this invention as the active component. The amount of oligonucleotide in each dose, and the administration schedule, can vary according to the corporal mass of the subject and stage in tumor progression. For example, for a human of about 70 kg who has an advanced, unresectable metastatic melanoma, a dose of about 1 mg of the oligonucleotide of this invention can be administered 3 times per week for about 10 weeks.

#### EXAMPLE 16

##### Induction of peripheral white blood cell IL-10 secretion and inhibition of IL-5 secretion by phosphorothioate non-CpG oligonucleotides of this invention

IL-10 is a potent suppressor of IgE production and induction and maintenance of epitope-specific T cell anergy is associated with increased IL-10. Fig 13 shows the stimulation of IL-10 secretion by the phosphorothioate non-CpG oligonucleotide IMT 504 and controls in PMBCs of several donors. As can be observed, IMT 504 is as active as the CpG positive control ODN 2006. Allergen-specific Th2 cells produce IL-5, which promotes the differentiation, activation and survival of eosinophils. Fig 14 shows that the phosphorothioate non-CpG oligonucleotide IMT 504 and the CpG positive control ODN 2006



are repressors of the production of IL-5 in several human PMBCs induced by IL-4 + Con A.

#### EXAMPLE 17

##### Treatment of subjects with allergic disease

5        Pharmaceutical formulations containing one or more of the immunostimulatory oligonucleotides of this invention, can be used to treat subjects against a variety of allergic diseases.

      In particular, a human with asthma can be treated by administration in the respiratory tract of an aerosolized pharmaceutical formulation containing the  
10       oligonucleotide of this invention as the active component. The amount of oligonucleotide in each dose, and the administration schedule, can vary according to the subject.

##### Oligonucleotide sequences which are outside the scope of the invention

15       The following oligonucleotide sequences (disclosed in the above listed prior art publications) are excluded from the scope of the immunostimulatory oligonucleotide claims; they might however be comprised within the scope of the composition and use claims:

20       ACTTCTCATAGTCCTTTGGTCCAG  
         TTGTTTTTTTGTGTTTTGTTTT  
         TTTTTTTTGTGTTTTGTTTT  
         TTTTGGGGTTTTGGGGTTTT  
         CTGGTCTTTCTGGTTTTTTCTGG  
25       CTGTGCTTTCTGTGTTTTTTCTGTG  
         GAACCTTCCATGCTGTT

GCTGAACCTTCCATGCTGTT  
TTTTTGTGTTTGTGGTTTGTGGTT  
TTGGTTTTTTTGGTTTTTTTTTGG  
ACATCTGGTTCTTACTTCAGG  
5 GGTTCCTTTTGGTCCTTGTCT  
GGTTCCTTTTGGTCCTTATCT  
TGGTCTTTTGGTCCTTGTCT  
TTCAGTTGTCTTGCTGCTTAGCTAA  
GGTTCCTTTTGGTCCTTGTCT  
10 TGGTGGTTTTGTGGTTTTGTGGTT  
TCCTCCTTTTGTCTTTTGTCTT  
CTGTGCTTTCTGTGTTTTCTGTG  
TGCTGCTTGTGCTTGTGCTT  
GGTTCCTTTTGGTCCTTGTCT  
15 CTGGTCTTTCTGGTTTTTTCTGG  
TNGTNGTTTTGTNGTTTTGTNGTT  
CTGTGCTTTCTGTGTTTTCTGTG  
GAACCTTCCATGCTGTT  
GCTGAACCTTCCATGCTGTT  
20 TTTTGTGTTTGTGGTTTTGTGGTT  
TTCAGTTGTCTTGCTGCTTAGCTAA  
TTGGTTTTTTTGGTTTTTTTTTGG  
ACATCTGGTTCTTACTTCAGG  
GGTTCCTTTTGGTCCTTATCT  
25 TGGTCTTTTGGTCCTTGTCT  
TTGTTGTTGTTGTTTGTGTTGTTG  
TCCATGTNGTTCCTGATGCT  
ACTTCTCATAGTCCCTTTGGTCCAG  
TTGTTTTTTTGTGTTTTTTGTTTT  
30 TTTTTTTTGTGTTTTTTGTTTT  
CTTAGGGGTGTTCTTAGGTGGTAACCCCTAGGGGTACACCTTCATTG  
GAAAACCTT  
TTAGTTGTTAGTTATTAGTT  
TTAGTTGTTAGTTTTTAGTT  
35 TTAGTTGTTAGTTCTTAGTT  
TTAGTTGTTAGTTGTTAGTT  
GGCTGAGCTGGCTTTTTCTTGTGG  
CTGGTCTTTCTGGTTTTTTCTGG  
CTCCTATTGGTGGTTTCCTAT  
40 CTCCTATTGGGTTTTTCCTAT  
CTCCTATTGGTGTTCCTAT  
CTCCTATTGGTGTTCCTAT  
CTCCTATTGGTGTTCCTAT  
CTCCTATTGTGGTTTCCTAT

CTCCTATTTTGGGTTTCCTAT  
TTGGAGTTTTTGGTGGGG  
TTGGAGTTTTTGGTTTTT  
CTGATGTGTTGAAGAACA  
5 TGCTGTATTTCTGGTACA  
GTTATCCCTGCTGT  
ACCTTGCTGTACTG  
CTGGCTTTTGGGTT  
TCTGTTGTGACTCAAG  
10

## CLAIMS

1. An immunostimulatory oligonucleotide having about 14 to 100 nucleotides, comprising a non-palindromic nucleic acid sequence motif having the following formula:



wherein  $X_1$  is C or T,  $X_2$  is C, T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are T; with the proviso when  $X_1$  is C,  $X_2$  is T, G or A and that, in the motif, a C does not precede a G.

2. An immunostimulatory oligonucleotide having about 14 to 100 nucleotides, comprising a non-palindromic nucleic acid sequence motif having the following formula:



wherein  $X_1$  is C,  $X_2$  is T, G or A,  $X_3$  is T or A,  $X_4$  is T, C or G,  $X_5$  is T, C or G,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T and wherein at least two of  $X_3$ ,  $X_4$ ,  $X_5$  and  $X_6$  are T; with the proviso and that, in the motif, a C does not precede a G.

3. The immunostimulatory oligonucleotide of claim 1, wherein  $X_1$  is C.
4. The immunostimulatory oligonucleotide of claims 1-2, wherein all of  $X_3$ ,  $X_4$ ,  $X_5$ ,  $X_6$  and  $X_8$  are T.
5. The immunostimulatory oligonucleotide of claims 1-4, wherein  $X_3X_4X_5X_6X_7X_8$  consists of TTTTGT.

6. The immunostimulatory oligonucleotide of claims 1-2 wherein  $X_1$  is C,  $X_3$  is T or A,  $X_4$  is T,  $X_5$  is T,  $X_6$  is T or G,  $X_7$  is G and  $X_8$  is T.
7. The immunostimulatory oligonucleotide of claim 1 having about 14 to 40 nucleotides.
- 5 8. The immunostimulatory oligonucleotide of claim 1 having a modified phosphate backbone.
9. The immunostimulatory oligonucleotide of claim 8 having a phosphate backbone modification on the 5' inter-nucleotide linkages.
10. The immunostimulatory oligonucleotide of claim 8 having a phosphate backbone modification on the 3' inter-nucleotide linkages.
- 10 11. The immunostimulatory oligonucleotide of claims 1-8 wherein at least one of the internucleotide linkages is a phosphorotioate linkage.
12. The immunostimulatory oligonucleotide of claims 1-11, having a sequence comprising:
 

$$5'-N X_1 X_2 X_3 X_4 X_5 X_6 X_7 X_8 N-3'$$

wherein N is an oligonucleotide composed of about 3-46 nucleotides, preferably of about 3-25 nucleotides.
13. The immunostimulatory oligonucleotide of claim 12, having a sequence comprising:
 

$$5'-NCX_2 TTTTGTN-3'$$

wherein  $X_2$  is T or G or C or A.

14. An immunostimulatory oligonucleotide which consists of a nucleotide sequence selected from:

- TCATCATT TTTGTCATT TTTGTCATT (SEQ ID N°2);
- TCTTCT TTTTGTCT TTTTGTCTTT (SEQ ID N°4);
- 5 TTGTTG TTTTGT TTTTGT TTTGTT (SEQ ID N°7);
- TCATTT TTTTGT TTTT TTTGTCATT (SEQ ID N°10);
- TCATTG TTTTGT TTTT TTTGTCATT (SEQ ID N°11);
- TCATTCT TTTGTTCT TTTT TTTGTCATT (SEQ ID N°12);
- TCATTAT TTTGTTAT TTTT TTTGTCATT (SEQ ID N°15);
- 10 TCATCCT TTTGTCCT TTTT TTTGTCATT (SEQ ID N°17);
- TCATCT TTTTGTCT TTTT TTTGTCATT (SEQ ID N°18);
- CATTTT GTTTTT TTTTTTTTTTTT (SEQ ID N°20);
- TTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°21);
- TTTTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°22);
- 15 TTTTTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°23);
- TTTTTTTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°24);
- TTTTTTTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°25);
- TTTTTTTCATTT GTTTTT TTTTTTTTTTTT (SEQ ID N°26);
- TTTTTTTCATTT GTTTT TTTTTTTTTTTT (SEQ ID N°27);

TTTTTTTTTTTTTTTCATTTTGT (SEQ ID N° 28);  
TTTTCATTTTGTCAATTTGTTTT (SEQ ID N°29);  
TCATCAATTTGTCAATTTGTCATT (SEQ ID N°30);  
ACATCATTTTGTCAATTTGTCATT (SEQ ID N°46);  
5 CCATCATTTTGTCAATTTGTCATT (SEQ ID N°47);  
GCATCATTTTGTCAATTTGTCATT (SEQ ID N°48);  
TAATCATTTTGTCAATTTGTCATT (SEQ ID N°49);  
TTATCATTTTGTCAATTTGTCATT (SEQ ID N°50);  
TGATCATTTTGTCAATTTGTCATT (SEQ ID N°51);  
10 TCCTCATTTTGTCAATTTGTCATT (SEQ ID N°52);  
TCTTCATTTTGTCAATTTGTCATT (SEQ ID N°53);  
TCAACATTTTGTCAATTTGTCATT (SEQ ID N°55);  
TCACCATTTTGTCAATTTGTCATT (SEQ ID N°56);  
TCAGCATTTTGTCAATTTGTCATT (SEQ ID N°57);  
15 TCATCATTTTGTCAATTTGTAATT (SEQ ID N°58);  
TCATCATTTTGTCAATTTGTTATT (SEQ ID N°59);  
TCATCATTTTGTCAATTTGTGATT (SEQ ID N°60);  
TCATCATTTTGTCAATTTGTCCTT (SEQ ID N°61);  
TCATCATTTTGTCAATTTGTCTTT (SEQ ID N°62);

- TCATCATTTTGTCAATTTTGTCAAT (SEQ ID N°64);
- TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°65);
- TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°66);
- TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°67);
- 5 TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°68);
- TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°69);
- TCATCAATTGGTCAATTGGTCATT (SEQ ID N°75);
- TCATCAACTGGTCAACTGGTCATT (SEQ ID N°77);
- TCATCATTTGTCAATTTGTCACT (SEQ ID N°84);
- 10 TCATCATTTGGTCAATTTGGTCATT (SEQ ID N°87);
- TCATCAATTGTCAATTTGTCACT (SEQ ID N°91);
- TCATCATCTTGTCACTTGTCACT (SEQ ID N°95);
- TCATCATGTTGTCACTTGTCACT (SEQ ID N°96).

15. An immunostimulatory oligonucleotide which is:

- 15 TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°2).

16. An immunostimulatory oligonucleotide which contains a nucleotide sequence selected from the group consisting of:

- TCATCATTTTGTCAATTTTGTCACT (SEQ ID N°2);
- TCTTCTTTTGTCTTTTGTCTTT (SEQ ID N°4);
- 20 TTGTTGTTTGTGTTTGTGTTT (SEQ ID N°7);



TCATTTTTTTGTTTTTTGTCATT (SEQ ID N°10);  
TCATTGTTTTGTTGTTTTGTCATT (SEQ ID N°11);  
TCATTCTTTTGTTCTTTTGTCATT (SEQ ID N°12);  
TCATTATTTTGTTATTTTGTCATT (SEQ ID N°15);  
5 TCATCCTTTTGTCCTTTTGTCATT (SEQ ID N°17);  
TCATCTTTTGTCCTTTTGTCATT (SEQ ID N°18);  
CATTTGTTTTTTTTTTTTTTTTTTT (SEQ ID N°20);  
TTCATTTTGTTTTTTTTTTTTTTTTT (SEQ ID N°21);  
TTTTCATTTTGTTTTTTTTTTTTTTT (SEQ ID N°22);  
10 TTTTTTCATTTTGTTTTTTTTTTTTT (SEQ ID N°23);  
TTTTTTTTTCATTTTGTTTTTTTTTT (SEQ ID N°24);  
TTTTTTTTTTTCATTTTGTTTTTTT (SEQ ID N°25);  
TTTTTTTTTTTTTCATTTTGTTTTT (SEQ ID N°26);  
TTTTTTTTTTTTTTTTTCATTTTGTTT (SEQ ID N°27);  
15 TTTTTTTTTTTTTTTTTTCATTTTGT (SEQ ID N°28);  
TTTTCATTTTGTCATTTTGTTTTT (SEQ ID N°29);  
TCATCAATTTGTCAATTTGTCATT (SEQ ID N°30);  
ACATCATTTTGTCATTTTGTCATT (SEQ ID N°46);  
CCATCATTTTGTCATTTTGTCATT (SEQ ID N°47);

GCATCATTTTGTGATTTTGTGATT (SEQ ID N°48);  
TAATCATTTTGTGATTTTGTGATT (SEQ ID N°49);  
TTATCATTTTGTGATTTTGTGATT (SEQ ID N°50);  
TGATCATTTTGTGATTTTGTGATT (SEQ ID N°51);  
5 TCCTCATTTTGTGATTTTGTGATT (SEQ ID N°52);  
TCTTCATTTTGTGATTTTGTGATT (SEQ ID N°53);  
TCAACATTTTGTGATTTTGTGATT (SEQ ID N°55);  
TCACCATTTTGTGATTTTGTGATT (SEQ ID N°56);  
TCAGCATTTTGTGATTTTGTGATT (SEQ ID N°57);  
10 TCATCATTTTGTGATTTTGTGAATT (SEQ ID N°58);  
TCATCATTTTGTGATTTTGTGTTATT (SEQ ID N°59);  
TCATCATTTTGTGATTTTGTGATT (SEQ ID N°60);  
TCATCATTTTGTGATTTTGTGCTT (SEQ ID N°61);  
TCATCATTTTGTGATTTTGTGCTT (SEQ ID N°62);  
15 TCATCATTTTGTGATTTTGTGAAT (SEQ ID N°64);  
TCATCATTTTGTGATTTTGTGACT (SEQ ID N°65);  
TCATCATTTTGTGATTTTGTGAGT (SEQ ID N°66);  
TCATCATTTTGTGATTTTGTGATA (SEQ ID N°67);  
TCATCATTTTGTGATTTTGTGATC (SEQ ID N°68);

TCATCATTTTGTCAATTTTGTGTCATG (SEQ ID N°69);

TCATCAATTGGTCAATTGGTTCATT (SEQ ID N°75);

TCATCAACTGGTCAACTGGTTCATT (SEQ ID N°77);

TCATCATTGTGTCATTGTGTCATT (SEQ ID N°84);

5 TCATCATTTGGTCAATTTGGTTCATT (SEQ ID N°87);

TCATCAATTTGTCAATTTGTGTCATT (SEQ ID N°91);

TCATCATCTTGTGTCATCTTGTGTCATT (SEQ ID N°95);

TCATCATGTTGTGTCATGTTGTGTCATT (SEQ ID N°96).

10 17. The immunostimulatory oligonucleotide of claims 1-16 wherein the immunostimulatory oligonucleotide is encapsulated in a slow release delivery vehicle.

18. The immunostimulatory oligonucleotide of claims 1-16 wherein the immunostimulatory oligonucleotide is included in a pharmaceutically acceptable carrier.

15 19. A pharmaceutical composition comprising an immunostimulatory oligonucleotide according to claims 1-16 and a pharmaceutically acceptable carrier.

20. A composition according to claim 19 wherein said immunostimulatory oligonucleotide is included in a plasmid.

20 21. A composition according to claim 19 further comprising an antigen.

22. A composition according to claim 21 wherein the antigen is selected from the group consisting of viruses, bacteria, fungi, parasites, tumoral cells, toxins, allergens, proteins, glycolipids and polysaccharides.

23. A composition according to claim 21 wherein the antigen is a viral antigen,  
5 a bacterial antigen, a human or animal tumoral cell and/or a fungal antigen.

24. A composition according to claim 21 wherein the antigen is encoded by a plasmid.

25. A composition according to claims 19-24 which is in liquid or lyophilized  
10 form.

26. Use of an immunostimulatory oligonucleotide according to claims 1-16 for the manufacture of a medicament for inducing B-cell activation; for treating, preventing or ameliorating an immune system disorder; for treating, preventing or ameliorating a tumoral disease; and/or for inducing  
15 plasmacytoid dendritic cell activation in a subject.


27. The use of claim 26, wherein the subject is a human.

28. The use of claim 26 wherein the oligonucleotide has been conjugated with one or more antigens.

29. The use of claim 26 wherein the oligonucleotide and the antigen are  
20 administered contemporaneously or simultaneously.

30. The use of claim 26 wherein the medicament is administered by intramuscular, oral, intranasal, anal, vaginal or transdermic route.

- 1/14 -




---

IMT 021 (SEQ. ID N° 1); IMT 023 (SEQ. ID N° 9). ODNs at 1.5 ug/ml


**FIGURE 1**

**FIGURE 2**



IMT 021 (SEQ. ID N° 1); IMT 023 (SEQ. ID N° 9). ODNs at 1.5 ug/ml


- 3/14 -

**FIGURE 3**

IMT 023 (SEQ. ID N° 9); IMT 504 (SEQ. ID N° 2); IMT 059 (SEQ. ID N° 98); IMT 037 (SEQ. ID N° 100); IMT 058 (SEQ. ID N° 99); IMT 034 (SEQ. ID N° 101); IMT 057 (SEQ. ID N° 102); IMT 028 (SEQ. ID N° 103); IMT 040 (SEQ. ID N° 104); IMT 046 (SEQ. ID N° 105); IMT 543 (SEQ. ID N° 36); IMT 544 (SEQ. ID N° 37); IMT 545 (SEQ. ID N° 38); IMT 546 (SEQ. ID N° 39); IMT 550 (SEQ. ID N° 43); IMT 608 (SEQ. ID N° 75). ODNs at 0.375 µg/ml

- 4/14 -

**FIGURE 4**




IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 053 (SEQ. ID N° 19); ODN(O): phosphorodiester ODN. ODNs at 3 x 30 ug/ml




- 5/14 -

**FIGURE 5**




IMT 022 (SEQ. ID N° 8); IMT 053 (SEQ. ID N° 19); IMT 173 (SEQ. ID N° 20); IMT 174 (SEQ. ID N° 21); IMT 175 (SEQ. ID N° 22); IMT 176 (SEQ. ID N° 23); IMT 177 (SEQ. ID N° 24); IMT 178 (SEQ. ID N° 25); IMT 179 (SEQ. ID N° 26); IMT 180 (SEQ. ID N° 27); IMT 181 (SEQ. ID N° 28); IMT 182 (SEQ. ID N° 29). ODNs at 0.375 ug/ml

- 6/14 -

**FIGURE 6**


IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2). ODNs at 1.5  $\mu$ g/ml.

- 7/14 -

**FIGURE 7**

IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2); IMT 506 (SEQ. ID N° 4). ODNs at 1.5 ug/ml


**FIGURE 8**



IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2); IMT 506 (SEQ. ID N° 4); ODN(O): phosphorodiester ODN. ODNs at 3 x 30 ug/ml

- 9/14 -


**FIGURE 9**



IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2); IMT 506 (SEQ. ID N° 4). ODNs at 0.375 ug/ml

- 10/14 -

**FIGURE 10**




IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2). ODNs at 1.5 ug/ml


- 11/14 -

**FIGURE 11**

**14 hs.**




**5 Days**



IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2). ODNs at 1.5 ug/ml


**FIGURE 12**



IMT 021 (SEQ. ID N° 1); IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2). ODNs at 1.5 ug/ml




- 13/14 -

**FIGURE 13**

IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2). ODNs at 1.5 ug/ml

**FIGURE 14**



IMT 022 (SEQ. ID N° 8); IMT 504 (SEQ. ID N° 2); ODNs at 1.5 ug/ml

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5 (i) APPLICANT: IMMUNOTECH SA
- (ii) TITLE OF INVENTION: Immunostimulatory oligonucleotides and  
uses thereof.
- (iii) NUMBER OF SEQUENCES: 74
- (iv) CORRESPONDENCE ADDRESS:
- 10 (A) ADDRESSEE: Dragotti & Associati  
(B) STREET: Galleria San Babila 4/C  
(C) CITY: Milano  
(D) STATE:  
(E) COUNTRY: Italy  
(F) ZIP: 20122
- 15 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM:  
(D) SOFTWARE:
- 20 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- (vii) PRIORITY APPLICATION DATA:

(A) APPLICATION NUMBER: CA-2388049

(B) FILING DATE: 2002-05-30

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Roberto Pistolesi

5 (B) REGISTRATION NUMBER: 94880

(C) REFERENCE/DOCKET NUMBER: 03mg06e

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: +39 02 799340

(B) TELEFAX: +39 02 784427

10

(2) INFORMATION FOR SEQ ID N°1 (IMT 021):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 1:

TGCTGCTTTTGTGCTTTTGTGCTT

20

(2) INFORMATION FOR SEQ ID N°2 (IMT 504):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

25 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 2:

5 TCATCATTTTGTTCATTTTGTTCATT

(2) INFORMATION FOR SEQ ID N°3 (IMT 505):

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

15 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 3:

TCCTCCTTTTGTCTTTTGTCTT

(2) INFORMATION FOR SEQ ID N°4 (IMT 506):

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 4:

TCTTCTTTTTGTCTTTTTGTCTTT

(2) INFORMATION FOR SEQ ID N°5 (IMT 501):

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 5:

TAGTAGTTTTGTAGTTTTGTAGTT

(2) INFORMATION FOR SEQ ID N°6 (IMT 502):

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 6:

TGGTGGTTTTGTGGTTTTGTGGTT

25 (2) INFORMATION FOR SEQ ID N°7 (IMT 503):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 7:

TTGTTGTTTGTGTTGTTTGTGTTGTT

10 (2) INFORMATION FOR SEQ ID N°8 (IMT 022):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 8:

TGCTGCAAAAGAGCAAAAGAGCAA

20 (2) INFORMATION FOR SEQ ID N°9 (IMT 023):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

25 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 9:

TGCTGCCCCCGCGCCCCGCGCCC

5

(2) INFORMATION FOR SEQ ID N°10 (IMT 197):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

10

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 10:

TCATTTTTTTGTTTTTTTGTTCATT

15

(2) INFORMATION FOR SEQ ID N°11 (IMT 198):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

20

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 11:

25

TCATTGTTTTGTTGTTTTGTTCATT



## (2) INFORMATION FOR SEQ ID N°12 (IMT 199):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

5 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 12:

10 TCATTCTTTTGTTCCTTTTGTCATT

## (2) INFORMATION FOR SEQ ID N°13 (IMT 509):

## (i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## 20 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 13:

AAAAAACTAAAAAAACTAAAAAA

## (2) INFORMATION FOR SEQ ID N°14 (IMT 531):

## 25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 14:

TCATAATTTTGTAATTTTGTTCATT

(2) INFORMATION FOR SEQ ID N°15 (IMT 532):

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 15:

TCATTATTTTGTTATTTTGTTCATT

20 (2) INFORMATION FOR SEQ ID N°16 (IMT 533):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 16:

TCATGATTTTGTGATTTTGTTCATT

5 (2) INFORMATION FOR SEQ ID N°17 (IMT 534):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 17:

TCATCCTTTTGTCTTTTGTTCATT

15

(2) INFORMATION FOR SEQ ID N°18 (IMT 535):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 18:

TCATCTTTTTGTCTTTTGTTCATT

25

(2) INFORMATION FOR SEQ ID N°19 (IMT 053):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

5 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 19:

TTTTTTTTTTTTTTTTTTTTTTTTTTT

10

## (2) INFORMATION FOR SEQ ID N°20 (IMT 173):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 20:

CATTTTGTTTTTTTTTTTTTTTTTTT

20

## (2) INFORMATION FOR SEQ ID N°21 (IMT 174):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

25 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 21:

5 TTCATTTTGTTTTTTTTTTTTTTT

(2) INFORMATION FOR SEQ ID N°22 (IMT 175):

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

15 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 22:

TTTTCATTTTGTTTTTTTTTTTTTT

(2) INFORMATION FOR SEQ ID N°23 (IMT 176):

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 23:

TTTTTTCATTTTGTTTTTTTTTTT

(2) INFORMATION FOR SEQ ID N°24 (IMT 177):

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 24:

TTTTTTTTTCATTTTGTTTTTTTTT

15 (2) INFORMATION FOR SEQ ID N°25 (IMT 178):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 25:

TTTTTTTTTTTCATTTTGTTTTTTTTT

25

(2) INFORMATION FOR SEQ ID N°26 (IMT 179):

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: DNA
- (iii) SEQUENCE DESCRIPTION: SEQ ID N° 26:

TTTTTTTTTTTTTCATTTTGTTTT

- 10 (2) INFORMATION FOR SEQ ID N°27(IMT 180):

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 15 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA
- (iii) SEQUENCE DESCRIPTION: SEQ ID N° 27:

TTTTTTTTTTTTTTTTTCATTTTGTTT

- 20 (2) INFORMATION FOR SEQ ID N°28 (IMT 181):

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- 25 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 28:

TTTTTTTTTTTTTTTTTCATTTTGT

5

(2) INFORMATION FOR SEQ ID N°29 (IMT 182):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 29:

TTTTCATTTTGTCATTTTGTTTTT

15

(2) INFORMATION FOR SEQ ID N°30 (IMT 537):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 30:

25 TCATCAATTTGTCAATTTGTCATT



## (2) INFORMATION FOR SEQ ID N°31 (IMT 538):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

5 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 31:

10 TCATCATATTGTCATATTGTCATT

## (2) INFORMATION FOR SEQ ID N°32 (IMT 539):

## (i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## 20 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 32:

TCATCATTATGTCATTATGTCATT

## (2) INFORMATION FOR SEQ ID N°33 (IMT 540):

## 25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5 (iii) SEQUENCE DESCRIPTION: SEQ ID N° 33:

TCATCATTTAGTCATTTAGTCATT

(2) INFORMATION FOR SEQ ID N°34 (IMT 541):

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 34:

TCATCATTTTATCATTTTATCATT

20 (2) INFORMATION FOR SEQ ID N°35 (IMT 542):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 35:

TCATCATTTTTTCATTTTTTCATT

5 (2) INFORMATION FOR SEQ ID N°36 (IMT 543):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 36:

TCATCATTTTCTCATTTTCTCATT

15 (2) INFORMATION FOR SEQ ID N°37 (IMT 544):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 37:

TCATCATTTTGACATTTTGACATT

25

(2) INFORMATION FOR SEQ ID N°38 (IMT 545):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 38:

TCATCATTTAGACATTTAGACATT

## (2) INFORMATION FOR SEQ ID N°39 (IMT 546):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA

## (iii) SEQUENCE DESCRIPTION: SEQ ID N° 39:

TCATCATTATGACATTATGACATT

## (2) INFORMATION FOR SEQ ID N°40 (IMT 547):

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 40:

TCATCATATTGACATATTGACATT

5

(2) INFORMATION FOR SEQ ID N°41 (IMT 548):

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

10 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) SEQUENCE DESCRIPTION: SEQ ID N° 41:

15 TCATCATTAAGACATTAAGACATT

(2) INFORMATION FOR SEQ ID N°42 (IMT 549):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 42:

25 TCATCATATAGACATATAGACATT

## (2) INFORMATION FOR SEQ ID N°43 (IMT 550):

## (A) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (B) MOLECULE TYPE: DNA

## 10 (C) SEQUENCE DESCRIPTION: SEQ ID N° 43:

TCATCATAATGACATAATGACATT

## (2) INFORMATION FOR SEQ ID N°44 (IMT 551):

## 15 (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## 20 (B) MOLECULE TYPE: DNA

## (C) SEQUENCE DESCRIPTION: SEQ ID N° 44:

TCATCATAAAGACATAAAGACATT

## 25 (2) INFORMATION FOR SEQ ID N°45 (IMT 552):

## (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 45:

TCATCAAAAAGACAAAAGACATT

10 (2) INFORMATION FOR SEQ ID N°46 (IMT 559):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 15 (D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 46:

ACATCATTTTGTTCATTTTGTTCATT

20 (2) INFORMATION FOR SEQ ID N°47 (IMT 560):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 25 (D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 47:

CCATCATTTTGTTCATTTTGTTCATT

5

(2) INFORMATION FOR SEQ ID N°48 (IMT 561):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

10

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 48:

15

GCATCATTTTGTTCATTTTGTTCATT

(2) INFORMATION FOR SEQ ID N°49 (IMT 562):

(A) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

25

(C) SEQUENCE DESCRIPTION: SEQ ID N° 49:

TAATCATTTTGTTCATTTTGTTCATT



## (2) INFORMATION FOR SEQ ID N°50 (IMT 563):

## (A) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (B) MOLECULE TYPE: DNA

## 10 (C) SEQUENCE DESCRIPTION: SEQ ID N° 50:

TTATCATTTTGTTCATTTTGTTCATT

## (2) INFORMATION FOR SEQ ID N°51 (IMT 564):

## 15 (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## 20 (B) MOLECULE TYPE: DNA

## (C) SEQUENCE DESCRIPTION: SEQ ID N° 51:

TGATCATTTTGTTCATTTTGTTCATT

## 25 (2) INFORMATION FOR SEQ ID N°52 (IMT 565):

## (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 52:

TCCTCATTTTGTGTCATTTTGTGTCATT

10 (2) INFORMATION FOR SEQ ID N°53 (IMT 566):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 53:

TCTTCATTTTGTGTCATTTTGTGTCATT

20

(2) INFORMATION FOR SEQ ID N°54 (IMT 567):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 54:

TCGTCATTTTGTTCATTTTGTTCATT

5

(2) INFORMATION FOR SEQ ID N°55 (IMT 568):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 55:

TCAACATTTTGTTCATTTTGTTCATT

15

(2) INFORMATION FOR SEQ ID N°56 (IMT 569):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 56:

25

TCACCATTTTGTTCATTTTGTTCATT

## (2) INFORMATION FOR SEQ ID N°57 (IMT 570):

## (A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

5 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

## (C) SEQUENCE DESCRIPTION: SEQ ID N° 57:

10 TCAGCATT TTTGTCATT TTTGTCATT

## (2) INFORMATION FOR SEQ ID N°58 (IMT 571):

## (A) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

20 (C) SEQUENCE DESCRIPTION: SEQ ID N° 58:

TCATCAT TTTTGTCA TTTTGTAA TT

## (2) INFORMATION FOR SEQ ID N°59 (IMT 572):

25 (A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

5 (C) SEQUENCE DESCRIPTION: SEQ ID N° 59:

TCATCATTTTGTTCATTTTGTATT

(2) INFORMATION FOR SEQ ID N°60 (IMT 573):

10 (A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15 (B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 60:

TCATCATTTTGTTCATTTTGTGATT

20 (2) INFORMATION FOR SEQ ID N°61 (IMT 574):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 61:

TCATCATTTTGTTCATTTTGCCTT

5 (2) INFORMATION FOR SEQ ID N°62 (IMT 575):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 62:

TCATCATTTTGTTCATTTTGTCTTT

15

(2) INFORMATION FOR SEQ ID N°63 (IMT 576):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 63:

TCATCATTTTGTTCATTTTGTGCGTT

25

(2) INFORMATION FOR SEQ ID N°64 (IMT 577):

## (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
5 (D) TOPOLOGY: linear

## (B) MOLECULE TYPE: DNA

## (C) SEQUENCE DESCRIPTION: SEQ ID N° 64:

TCATCATTTTGTCAATTTTGTCAAT

10

## (2) INFORMATION FOR SEQ ID N°65 (IMT 578):

## (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
15 (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (B) MOLECULE TYPE: DNA

## (C) SEQUENCE DESCRIPTION: SEQ ID N° 65:

TCATCATTTTGTCAATTTTGTCACT

20

## (2) INFORMATION FOR SEQ ID N°66 (IMT 579):

## (A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs  
25 (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 66:

TCATCATTTTGTTCATTTTGTTCAGT

5

(2) INFORMATION FOR SEQ ID N°67 (IMT 580):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

10

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 67:

15

TCATCATTTTGTTCATTTTGTTCATA

(2) INFORMATION FOR SEQ ID N°68 (IMT 581):

(A) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

25

(C) SEQUENCE DESCRIPTION: SEQ ID N° 68:



TCATCATTTTGTTCATTTTGTTCATC

(2) INFORMATION FOR SEQ ID N°69 (IMT 582):

5 (A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10 (B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 69:

TCATCATTTTGTTCATTTTGTTCATG

15 (2) INFORMATION FOR SEQ ID N°70 (IMT 187):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 70:

TTTTCATTTTGT

25

(2) INFORMATION FOR SEQ ID N°71 (IMT 188):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5 (B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 71:

TTTTCATTTTGTTTTT

10 (2) INFORMATION FOR SEQ ID N°72 (IMT 189):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 72:

TTTTCATTTTGTTTTTTTTT

20

(2) INFORMATION FOR SEQ ID N°73 (IMT 179):

(A) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 73:

TTTTTTTTTTTCATTTTGTTTTT

5

(2) INFORMATION FOR SEQ ID N°74 (IMT 191):

(A) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear


(B) MOLECULE TYPE: DNA

(C) SEQUENCE DESCRIPTION: SEQ ID N° 74:

TTTTCATTTTGTTTTTTTTTTTTTTT

15

20



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**